

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 118 669 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
25.07.2001 Bulletin 2001/30

(51) Int Cl.⁷: **C12N 15/82, A01H 5/00,
C07K 1/00**

(21) Application number: **00310997.2**

(22) Date of filing: **08.12.2000**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **17.12.1999 EP 99310188**

(71) Applicants:
• **UNILEVER PLC**
London EC4P 4BQ (GB)
Designated Contracting States:
GB IE
• **UNILEVER N.V.**
3013 AL Rotterdam (NL)
Designated Contracting States:
BE CH DE DK ES FR GR IT LI NL PT SE AT

(72) Inventors:
• **Frenken, Leo Gerard Joseph**
3130 AC Vlaardingen (NL)
• **Van der Logt, Cornelis Paul Erik**
3130 Vlaardingen (NL)
• **Jobling, Stephen Alan**
Sharnbrook, Bedford, MK44 1LQ (GB)
• **Teh, Yih-Miin**
Sharnbrook, Bedford, MK44 1LQ (GB)

(74) Representative: **Van Velzen, Maaïke Mathilde et al**
Unilever N.V.
Patent Department
Postbus 137
3130 AC Vlaardingen (NL)

(54) **Production of camelid antibodies in plants**

(57) A method for modifying a plant to produce antibodies or active fragments or derivatives thereof in a desired cellular compartment comprising introducing in-

to a plant a DNA sequence encoding a heavy chain immunoglobulin or an active fragment or derivative thereof or a sequence encoding a protein functionally equivalent thereto and plants so modified are disclosed.

EP 1 118 669 A2

DescriptionFIELD OF THE INVENTION

5 **[0001]** The present invention is in the field of applied biotechnology and relates in particular to an economic way of producing antibodies, or more particularly fragments thereof, in plants.

BACKGROUND OF THE INVENTION

10 **[0002]** The production of antibodies in microbial or plant systems can be advantageous for a number of applications.
[0003] Firstly microbial or plant sources can be used as bioreactors. In this situation the antibodies and/or their derivatives are produced on a large scale in modified organisms. The use of such 'bioreactors', especially plants has the advantage that scale up to produce large quantities e.g. more than 100 kg or even more than 1000 kg of antibodies is relatively easy and does not require significant investments in harvesting or processing equipment.

15 **[0004]** Alternatively antibodies and/or their derivatives can be produced in plants with the aim of reprogramming the plant metabolism or to improve defence mechanisms of said plant. The antibodies to be produced are chosen such that they target specific enzymes to modulate their activity and/or to provide the plant with protection against pathogens such as parasites and viruses.

[0005] Several attempts have been made to express antibodies or fragments thereof in micro-organisms or plants.

20 **[0006]** Evidence from the literature suggests that where it is desired to express antibodies in microbial systems, the most favourable results are obtained using smaller fragments. Whole antibodies and larger antibody fragments, such as Fab fragments, are very difficult if not impossible to produce. Smaller molecules such as single chain antibody fragments (scFv) are somewhat easier. Generally, however the best results are obtained if even smaller molecules such as heavy chain variable fragments (HCV) are produced (see, for example, WO 94/25591 (Unilever)).

25 **[0007]** In plant host systems, both large molecules, such as complete murine antibodies and smaller murine antibody fragments such as single chain Fv (scFv) fragments are capable of being expressed.

[0008] The production of functional complete murine antibodies in plants was first reported by Hiatt et al, Nature, 342, 76-78 (1989). Subsequent reports of the functional expression of murine monoclonal antibodies in plants include Düring et al, Plant Molecular Biology, 15, 281-293, (1990) and Ma et al, European Journal of Immunology, 24, 131-138, 30 (1994).

[0009] Transgenic tobacco plants expressing a synthetic gene encoding an antigen binding single chain Fv protein (scFv) and which produce functional scFv protein have been described by Owen et al, Biotechnology, 10, 790-794, (1992). The expression of functional (that is having antigen-binding activity) scFv protein in transgenic plants has also been described in other reports in the literature, see, for example, Tavlaoraki et al, Nature, 366, 469-472 (1993).

35 **[0010]** Given that the production of functional complete antibodies requires the correct assembly, via covalent and non-covalent interactions, of both the antibody heavy and light chains, it might have been expected that expression of smaller antibody fragments, with their less stringent assembly requirements, would be advantageous. It has however been reported that generally, in practice, better yields are achieved with plants transformed with complete murine antibodies rather than small fragments (Ma et al, Science, 268, 716-719 (1995))

40 **[0011]** Furthermore, it is the experience of the present inventors that the expression of genes encoding scFv proteins in plants is not reliably reproducible and hence such a process would not readily lend itself to large scale production. Moreover, expression of genes encoding scFv molecules in plants can have an undesirable effect on plant cell morphology.

45 **[0012]** Benevenuto et al, Plant Molecular Biology, 17, 865-874 (1991) describe attempts to express smaller isolated murine heavy chain variable domain antibody fragments in plants. Successful expression is reported but there is no indication that binding affinity is retained.

[0013] For many purposes, it is important to be able to target antibodies or fragments thereof to specific cellular compartments. For example, where maximisation of yield is desired, targeting to the ER or the apoplastic space can help by providing a less reducing, more stable environment than is present in the cytoplasm. On the other hand, where 50 it is desired to interfere with the activity of specific enzymes, in order to engineer metabolic changes in a plant, it is necessary to ensure that the antibodies are directed to the particular compartment in which the enzyme is active.

[0014] In order to direct the expression of the desired protein to a specific cellular compartment within a plant cell, the antibody proteins are fused to specific targeting sequences (peptides) where required. Expression in the chloroplast requires an N-terminal chloroplast targeting sequence (generally termed transit peptide) which is cleaved off upon 55 entry into the chloroplast (reviewed in Keegstra & Cline (2000) Plant Cell 11, 557-570). There is little sequence homology among these transit peptides except that they are often rich in hydroxylated amino acids (eg serine), are deficient in acidic amino acids and they can vary in length from 30 to >100 amino acids. Various transit peptides fused to heterologous proteins have been used successfully to direct the protein to the chloroplast however there are no clear

rules that will guarantee correct targeting and cleavage of the fusion protein. Some groups have used an exact fusion of the transit peptide to the heterologous protein; for example the transit peptide of the small subunit of rubisco from petunia has been used in this way (Fray et al (2000) Nature Biotech 17, 1017-1020). Others have used transit peptides with two amino acids of the mature protein, eg potato granule-bound starch synthase (GBSS) (Kortsee et al (1996) Plant Journal 10, 83-90), and some have used as many as 23 amino acids of the mature protein eg the small subunit of rubisco from pea (Nawrath et al (1994) Proc. Natl. Acad. Sci USA 91, 12760-12764). Another group used the transit peptide of the Arabidopsis small subunit of rubisco plus 24 amino acids of the mature protein followed by a repeat of six amino acids before the cleavage site and an additional two amino acids of the mature rubisco protein (Wong et al (1992) Plant Mol. Biol. 20, 81-93).

[0015] Proteins that enter the secretory system have N-terminal targeting peptides generally known as "signal peptides or sequences" which direct them initially to the endoplasmic reticulum (ER) whence, depending on additional sequences, the proteins are sorted either to vacuoles or are secreted from the cell. In addition, proteins which enter the ER and have a carboxy terminal peptide KDEL are retained within the ER membrane system. For location in the cytoplasm, no targeting sequence is required.

[0016] The targeting of classical conventional antibodies (ie IgG, scFv etc) to various cellular compartments of plants has been reviewed in Conrad and Fielder (1998) Plant Molecular Biology 38, 101-109. Most commonly the homologous signal sequence of the murine IgG (e.g. CEA66E3) has been used to direct expression of the antibody via the secretory pathway, although the signal sequences from plants (PR1a, legumin B4, 2S storage protein) and even bacteria (pelB) and yeast have been used successfully. Generally the highest expression levels were obtained by the use of a signal peptide and C-terminal KDEL ER retention signal. There are no reports of successful targeting of antibodies to chloroplasts using a transit peptide as the targeting sequence. However, Düring et al (1990) Plant Mol. Biol. 15, 281-293, reported that when the barley α -amylase signal peptide was fused to the light and heavy chains of a monoclonal antibody, there was some apparent aberrant localisation of antibody within the chloroplast although the majority of the labelling was in the ER. The authors had no explanation for this observation and were unsure whether import into the chloroplast had actually occurred or if there were other reasons for the specific labelling. Interestingly, no labelling could be detected in the expected location of the apoplastic space or in the golgi apparatus or secretory vesicles.

[0017] WO 94/4678 (Casterman et al) describes the isolation of immunoglobulins from Camelids. These immunoglobulins, hereinafter "heavy chain immunoglobulins" have a characteristic distinct structure in that they are naturally devoid of light chains, with the antigen-binding sites being contained entirely within the heavy chain variable domains. This in turn leads to a characteristic structure for the heavy chains of these immunoglobulins, because the variable domain has no interaction sites for a light chain and a complete antigen-binding site has to be formed with no contribution from the hypervariable loops of a light chain variable domain. The heavy chain variable fragments (HCV) of these immunoglobulins are often referred to as VHH fragments. Such immunoglobulins or fragments thereof, show the functional properties of conventional, four chain, immunoglobulins but by virtue of their simplified structure offer advantages in preparation and use.

[0018] There are no reports in the literature of correctly targeted expression in plants of heavy chain immunoglobulins or fragments thereof. A report has appeared in a PhD thesis (Vu, 1999, Vrije Universiteit, Brussels, Belgium) of expression of a heavy chain variable domain in fusion with a chloroplast targeting peptide in *Nicotiana tabacum*. However, targeting to the chloroplast appears to have failed and the subcellular localisation of the antibody fragments (which remained as a fusion protein) was not established.

[0019] There remains a continuing need for the development of improved methods for the production of antibodies and fragments thereof in plants, with the aim of providing methods suitable for economical large scale production for protecting plants from pathogens and for modulating their metabolism.

SUMMARY OF THE INVENTION

[0020] Accordingly, in a first aspect the invention provides a method for modifying a plant to produce an antibody or an active fragment or derivative thereof in a desired cellular compartment, comprising introducing into a plant a DNA sequence encoding a heavy chain immunoglobulin or an active fragment or derivative thereof, or a sequence encoding a protein functionally equivalent thereto, said DNA sequence being operably linked to one or more promoters and provided, as appropriate, with an additional sequence encoding a peptide sequence capable of targeting said antibody or fragment or derivative thereof to said desired cellular compartment.

[0021] Also provided is a method for preparing a heavy chain immunoglobulin or an active fragment or derivative thereof comprising extracting said immunoglobulin or fragment or derivative thereof from a plant modified according to the first aspect of the invention.

[0022] In an alternative aspect, the invention provides the use of a DNA sequence encoding a heavy chain immunoglobulin or an active fragment or derivative thereof, or a sequence encoding a protein functionally equivalent thereto, to cause a plant to produce said antibody or fragment or derivative thereof or protein functionally equivalent thereto in

a desired cellular compartment.

[0023] Also provided is a modified plant having, in a desired cellular compartment, enhanced levels of heavy chain immunoglobulins or active fragments or derivatives thereof, or proteins functionally equivalent thereto, particularly compared to equivalent but unmodified plants.

[0024] Seeds, fruits and progeny of such plants and hybrids are included within the invention.

[0025] Food products such as sauces, dressings, tomato products such ketchups, meals, juices and soups, comprising a plant or part thereof according to the invention are also provided.

[0026] Also provided are skin and hair protective compositions and pharmaceutical compositions comprising a plant or part thereof according to the invention.

[0027] As used herein, "plant" means a whole plant or part thereof, or a plant cell or group of plant cells, or an extract thereof. The invention is particularly directed at transforming whole plants and the use of the whole plant or significant parts thereof.

[0028] The term "antibody" refers to an immunoglobulin which may be derived from natural sources or synthetically produced, in whole or in part. An "antibody fragment", alternatively an "active fragment", is a portion of a complete antibody or immunoglobulin which retains the ability to exhibit at least part, and preferably all, of the antigen binding activity.

[0029] A "heavy chain immunoglobulin" is an immunoglobulin naturally devoid of any variable light chain domains but which is capable of specifically combining with an antigen. The antigen-binding capacity and specificity is located exclusively in the immunoglobulin heavy chains, more specifically in the heavy chain variable domains.

[0030] The "sequence encoding the heavy chain immunoglobulin or an active fragment or derivative thereof" encompasses a genomic or cDNA clone or a sequence which in proper reading frame encodes an amino acid sequence which is functionally equivalent to the amino acid sequence of the protein encoded by the genomic or cDNA clone. By "functionally equivalent" is meant any protein or fragment or derivative thereof which has the same or similar antigen-binding properties, said antigen-binding capacity being located in a single binding domain. It should be understood, however, that isolated VH domains of conventional antibodies are not included within the scope of the invention. Similarity in functionality can be evaluated by routine screening assays, for example, by assaying the binding affinity of the immunoglobulins produced upon expression in plants.

[0031] A "functionally equivalent" derivative may be characterised by an insertion, deletion or substitution of one or more amino acid residues in the sequence of the heavy chain immunoglobulin or active fragment thereof and includes within its scope fusion molecules. Such derivatives may readily be made by using conventional techniques well known in the art such as peptide synthesis techniques or recombinant DNA manipulation of a gene encoding a heavy chain immunoglobulin by site directed mutagenesis.

[0032] It will be appreciated that immunoglobulins or fragments or derivatives thereof modified to enable them to function as binding domains in the same way as immunoglobulins naturally devoid of light chain domains ("heavy chain immunoglobulins") may also suitably be used according to the invention.

[0033] A "gene" is a DNA sequence encoding a protein, including modified or synthetic DNA sequences or naturally occurring sequences encoding a protein, and excluding the 5' sequence which drives the initiation of transcription.

[0034] "Operably linked to one or more promoters" means the gene, or DNA sequence, is positioned or connected to the promoter in such a way to ensure its functioning. The promoter is any sequence sufficient to allow the DNA to be transcribed. After the gene and promoter sequences are joined, upon activation of the promoter, the gene will be expressed.

[0035] An "equivalent, unmodified" plant is a plant which has a substantially identical genotype to a modified plant of the invention excepting the introduced sequences present in the plant of the invention.

[0036] A "construct" is a polynucleotide comprising nucleic acid sequences not normally associated in nature.

[0037] A "cellular compartment" is a volume or structure separated from the remainder of the cell by at least one membrane. For the purposes of the invention, this also includes the apoplastic space between cells. A "targeting peptide" or "transit peptide" is a polypeptide sequence which, when appended to the appropriate end of a protein sequence is capable of directing the import of said protein into a specific cellular compartment.

BRIEF DESCRIPTION OF THE DRAWINGS

[0038] The present invention may be more fully understood by reference to the following description, when read together with the accompanying drawings in which:

Figure 1 shows the nucleotide sequence (SEQ. ID. No. 1) and corresponding protein sequence (SEQ. ID. No. 2) of a fragment encoding the heavy chain variable domain of an anti-RR6 antibody (denoted HCV33) from a llama, with an attached peptide linker group (denoted myc) (RR6 is an azo-dye, available from ICI; myc is a peptide comprising the sequence EQKLISEEDLN, SEQ. ID. NO. 3).

- Figure 2 shows a specificity ELISA assay (A) and SDS-PAGE analysis (B) of whole cell extract of a tobacco plant transformed (Example 4) with the plant expression vector pPV.8 - HCV33-myc (of Example 1). The positive control was at a concentration of 250ng/ml. For the ELISA assay, 50µl of plant extract was used. Total protein concentration in the plant extract was 2mg/ml.
- Figure 3 shows a specificity ELISA assay (A) and SDS-PAGE analysis (B) of whole cell extracts of a tobacco plant transformed (Example 5) with the expression vector pPV.8-GBSS-HCV33-myc (of Example 1). The positive control was at a concentration of 20µg/ml. Total protein concentration in the plant extract was 1-2mg/ml.
- Figure 4 shows the results of immuno-histology studies, as reported in Example 5, on a tobacco plant transformed with the expression vector pPV.8-GBSS-HCV33-myc (of Example 1), confirming that HCV33 fragment is localised in the chloroplasts.
- Figure 5 shows the amount of active antibody present in fruit columella extracts of a tomato plant transformed (Example 5) with the plant expression vector pPV.8-GBSS-HCV33-myc (of Example 1), expressed as % of total soluble protein, as determined via direct binding ELISA.
- Figure 6 shows the nucleotide sequence (SEQ. ID. No. 4) and corresponding protein sequence (SEQ. ID. No. 5) of a fragment encoding the heavy chain variable domain of an anti-potato SBEII antibody (denoted Clonel) with attached peptide linker groups denoted His6 (His-His-His-His-His-His) and myc (as described in Figure 1). (Potato SBEII is a starch branching enzyme also denoted SBE A; Jobling et. al., Plant Journal, 18, 163-171 1999).
- Figure 7 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 5) with the plant expression vector pSJ35-CERV-GBSS-Clonel-His6-myc (of Example 1), expressed as % of total soluble protein, as determined via direct binding ELISA.
- Figure 8 shows the results of immuno-histology studies as reported in Example 6 on a tobacco plant transformed with the expression vector pSJ.34-2x355-GBSS-scFv3299-hydrophilIII (of Example 1). It is clear that plastid formation/function is disrupted.
- Figure 9 shows the amount of active antibody present in leaf extract of tobacco plants transformed (Example 7) with the plant expression vector pPV.8-PRIa-HCV33-myc (of Example 1), expressed as % of total soluble protein, as determined via direct binding ELISA.
- Figure 10 shows a specificity ELISA assay (A) and SDS-PAGE/western blot analysis (B) of whole cell extracts of tobacco plants transformed (Example 8) with two plant expression vectors, namely pPV.8-PRIa-HCV33-myc-SEKDEL and pPV.8-GBSS-HCV33-myc-SEKDEL (of Example 1). Total protein concentration in the plant extract was 1-2mg/ml.
- Figure 11 shows the results of immuno-histology studies as reported in Example 8 on a tobacco plant transformed with the expression vector pPV.8-PRIa-HCV33-myc-SEKDEL (of Example 1) confirming that the HCV33 material, directed to the secretory pathway using the PRIa leader sequence, and retained in the endoplasmic reticulum by the C-terminal tag SEKDEL.
- Figure 12 shows the nucleotide sequence (SEQ. ID. No. 6) and corresponding protein sequence (SEQ. ID. No. 7) of a fragment encoding the heavy chain variable domain of an anti-potato SBEII antibody (denoted Clone46) with attached peptide linker groups denoted His6 and myc (as described in Figure 1).
- Figure 13 shows that the SBE (starch branching enzyme) activity of whole leaf extracts of 30 tobacco plants independently transformed (Example 9) with the plant expression vector pSJ.35-CERV-GBSS-Clone46-His6-myc (of Example 1) is significantly less than that of plants transformed with pSJ.35-CERV-GBSS-HCV33-myc (of Example 1).
- Figure 14 shows the nucleotide sequence (SEQ. ID. No. 8) and corresponding protein sequence (SEQ. ID. No. 9) of a fragment encoding the heavy chain variable domain of an anti-GUS antibody (denoted Clone18) with attached peptide linker groups denoted His6 and myc (as described in Figure 1). (GUS is β-glucuro-

nidase, available from Sigma.)

Figure 15 shows the nucleotide sequence (SEQ. ID. No. 10) and corresponding protein sequence (SEQ. ID. No. 11) of a fragment encoding the heavy chain variable domain of an anti-GUS antibody (denoted Clone18) with an endoplasmic retention signal (SEKDEL) attached peptide linker groups denoted His6 and myc (as described in Figure 1).

Figure 16 shows that the GUS (β -glucuronidase) activity of whole leaf extracts of 30 pCJ.102 (of Example 1) transformed tobacco plants independently retransformed (Example 10) with the plant expression vector pSJ.35-CERV-Clone18-His6-myc-SEKDEL (of Example 1) is significantly less than that of control pCJ.102 plants.

Figure 17 shows (A) X-glu staining of pollen from a tobacco plant transformed (Example 11) with pSJ.35-Lat52-HCV33-myc-SEKDEL (of Example 1), and (B) X-glu staining of pollen from a tobacco plant transformed (Example 11) with pSJ.35-Lat52-Clone18-His6-myc-SEKDEL (of Example 1).

Figure 18 shows the nucleotide sequence (SEQ. ID. No. 54) and corresponding protein sequence (SEQ. ID. No. 55) of a fragment encoding the heavy chain variable domain of an anti-RR6 antibody (denoted HCV33) linked to the non-classical heavy chain constant regions (denoted hinge-CH2-CH3).

Figure 19 shows the nucleotide sequence (SEQ. ID. No. 56) and corresponding protein sequence (SEQ. ID. No. 57) of a fragment encoding the heavy chain variable domain of an anti-RR6 antibody (denoted HCV33) linked to the non-classical heavy chain constant regions (denoted Hinge-CH2-CH3) with an endoplasmic retention signal (SEKDEL).

Figure 20 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 12) with the plant expression vector pPV.8-HCV33-IgG2b (of Example 12), expressed as % of total soluble protein as determined via direct binding ELISA.

Figure 21 shows the amount of active antibody present in whole leaf, stem and root extract of three tobacco plants (from Figure 20) analysed 6 weeks post germination, expressed as % of total soluble protein as determined via direct binding ELISA (of Example 12).

Figure 22 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 12) with the plant expression vector pPV.8-GBSS-HCV33-IgG2b (of Example 12), expressed as % of total soluble protein as determined via direct binding ELISA.

Figure 23 shows the amount of active antibody present in whole leaf, stem and root extract of three tobacco plants (from Figure 22) analysed 6 weeks post germination, expressed as % of total soluble protein as determined via direct binding ELISA (of Example 12).

Figure 24 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 12) with the plant expression vector pPV.8-PR1a-HCV33-IgG2b (of Example 12), expressed as % of total soluble protein as determined via direct binding ELISA.

Figure 25 shows the amount of active antibody present in whole leaf, stem and root extract of three tobacco plants (from Figure 24) analysed 6 weeks post germination, expressed as % of total soluble protein as determined via direct binding ELISA (of Example 12).

Figure 26 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 12) with the plant expression vector pPV.8-PR1a-HCV33-IgG2b-SEKDEL (of Example 12), expressed as % of total soluble protein as determined via direct binding ELISA.

Figure 27 shows the amount of active antibody present in whole leaf, stem and root extract of three tobacco plants (from Figure 26) analysed 6 weeks post germination, expressed as % of total soluble protein as determined via direct binding ELISA (of Example 12).

Figure 28 shows the nucleotide sequence (SEQ. ID. No. 66) and corresponding protein sequence (SEQ. ID. No.

67) of a fragment encoding the heavy chain variable domain of an anti-potato SBEII antibody (denoted Clone68) with attached peptide linker groups denoted His6 and myc (as described in Figure 1).

Figure 29 shows the amount of active antibody present in whole leaf extract of a tobacco plant transformed (Example 13) with the plant expression vector pSJ35-CERV-GBSS-Clone68-His6-myc (of Example 13) expressed as OD reading as compared to the purified and quantified *E. coli* clone68 standard.

DETAILED DESCRIPTION OF THE INVENTION

[0039] The present invention is based on the finding that immunologically active immunoglobulins, or active fragments or derivatives thereof, which are devoid of light chain variable domains, can advantageously be expressed at high level and successfully targeted to specific cellular locations in plant cells, with retention of binding activity and no adverse effects on plant growth or morphology.

[0040] In particular, it has been found that the method of the invention can be used to produce significant amounts of whole heavy chain immunoglobulins, with retention of binding activity, in the major compartments of the plant cell, namely plastid, cytoplasm, apoplast and endoplasmic reticulum. Selfing of transformed, antibody-expressing plants produced a new generation of plants, many of which displayed an ability to accumulate higher levels of antibody than the parent plants. The amounts of antibody were found to be markedly greater than those typically achieved when conventional antibodies are expressed in plants. The accumulation of high levels of functional antibody in the cytoplasm was a particularly unexpected result, since conventional antibodies have generally been found to be produced only in very small amounts in this location, probably owing to the problem of forming the disulphide bonds needed for structural stability in a reducing environment.

[0041] The method of the invention was similarly found to be capable of producing VHH fragments in significant amounts in these different compartments. This was the case for several different VHH fragments, in contrast to the situation with fragments based on conventional antibodies, such as scFv, where the results have been found to be very unpredictable. Importantly, it was found that the accumulation of VHH fragments in any compartment of the cell resulted in no discernible deleterious effect on the plant; again this contrasts with results obtained with scFv fragments, where expression levels were found to be unstable and accumulation of the fragment in the plastids was found to affect adversely the structure and function of these organelles.

[0042] By means of the invention, antibodies and active fragments thereof, having specificity for a target antigen may conveniently be prepared by a method which is readily amenable to industrial-scale production. Further, the ability to target the antibodies to any chosen cell compartment makes it possible to use the method of the invention to modulate the activity of specific enzymes irrespective of where in the cell the enzyme is found.

[0043] In order to modify the function of an intracellular protein, the antibody or fragment or derivative thereof must be present and active in those compartments of the host plant cell where the target protein is active. In cases where this is the cytoplasm, the reducing environment inhibits the formation of disulphide links in the antibody and so generally antibodies would not be expected to be active in the cytoplasm. Expression of classical antibodies, or fragments thereof, in the cytoplasm does not generally give rise to functional activity, for example. Similarly, expression of scFv fragments in plastids generally leads to unstable deformed plastids. The targeted expression of active antibodies or fragments or derivatives thereof in compartments of the host plant cell, especially in the cytoplasm, therefore represents a particular advantage of the method of the invention.

[0044] The present inventors have found that even when the levels of expression of heavy chain immunoglobulin or fragment or derivative thereof in the plant cell are low or undetectable by conventional protein chemical methods such as in the case of expression in the cytoplasm, a measurable effect on enzyme activity may still be achieved. The ability to modulate plant enzyme activity without accumulating antibodies in large amounts may be beneficial in some circumstances, for example, in minimising potential deleterious effects on the growth and health of the plant.

[0045] The heavy chain immunoglobulins (or fragments or derivatives thereof) to be expressed in accordance to the invention are, or are derived from, immunoglobulins that are naturally devoid of light chains, their antigen binding sites being contained entirely within the heavy chain variable domains. Especially preferred are immunoglobulins or fragments or derivatives thereof which are obtainable from Camelids as described, for example, in WO 94/4678 above. Heavy chain variable domains of immunoglobulins naturally devoid of light chains are particularly preferred for use in the present invention.

[0046] Preferred immunoglobulins for use in the invention are obtainable from *Camelids* especially from *Lamas* (for example *Lama glama*, *Lama vicugna* or *Lama Paccos*) or from *Camelus* (for example *Camelus dromedarius* or *Camelus bactrianus*).

[0047] Conveniently, a functionally equivalent protein shows at least 50% similarity to the respective amino acid sequence, preferably at least 70% similarity, more preferably at least 80%, most preferably 90-100% similarity to the respective amino acid sequence as determined by techniques well known in the art.

[0048] The invention also provides for the expression of complex derivative molecules comprising an immunoglobulin or fragment thereof as described above, associated or otherwise connected to one or more other molecules, for example, an enzyme or a further heavy chain variable domain.

[0049] The invention provides specifically for the targeting of expressed antibodies, or fragment or derivatives thereof, to specific cellular compartments. Where the desired compartment is the cytoplasm, this is the default location when no specific targeting sequence is appended to the antibody. For targeting to other compartments, sequences encoding appropriate targeting signals are fused to the sequence encoding the antibody.

[0050] Where it is desired to target an antibody to the chloroplast, a sequence encoding an appropriate transit peptide is provided, such that said peptide will be fused at the N-terminus of the expressed antibody and subsequently removed upon passage into the organelle. Any chloroplast-specific transit peptide may be used together, optionally, with a few (1-3) residues from the N-terminus of the mature protein to which said peptide is fused in nature. A number of such sequences, proven to be capable of directing successful targeting of heterologous proteins, are well known to those skilled in the art. Particularly preferred is the GBSS transit peptide sequence from potato (van der Leij et al (1991) Mol. Gen. Genet., 228, 240-248).

[0051] Where it is desired to direct secretion of an antibody into the apoplastic space, a sequence encoding an appropriate signal peptide is provided, such that said peptide will be fused at the N-terminus of the expressed antibody and subsequently removed upon passage into the endoplasmic reticulum, en route to secretion. Any signal peptide may be used, including a signal peptide that would naturally direct secretion of an antibody, or other signal peptides derived from animal, plant or microbial secreted or ER-localised proteins. Many such sequences are well known to those skilled in the art. Particularly preferred is the PR1a leader sequence (Firek et al; Plant Mol. Biol., 23 861-870 (1993)).

[0052] Where it is desired to direct an antibody to the endoplasmic reticulum, and to retain it there, a signal sequence is provided, as above, and additionally an ER-retention signal is provided. Conveniently, this takes the form of a C-terminal polypeptide extension comprising the sequence Lys-Asp-Glu-Leu (KDEL). Alternatively, the KDEL sequence may be introduced by mutation of the residues at the C-terminal end of the antibody, or fragment of derivative thereof.

[0053] It will be appreciated that the invention extends to any plant which is amenable to transformation. Suitable plants include tobacco, peas, potatoes, spinach, tomato and tea.

[0054] The DNA sequences of interest will preferably be operably linked (that is, positioned to ensure the functioning of) to one or more suitable promoters which allow the DNA to be transcribed. Suitable promoters, which may be homologous or heterologous to the gene, useful for expression in plants are well known in art, as described, for example, in Weising et al, (1988), Ann. Rev. Genetics, 22, 421-477). Promoters for use according to the invention may be inducible, constitutive or tissue-specific or have various combinations of such characteristics. Useful promoters include, but are not limited to constitutive promoters such as carnation etched ring virus (CERV), cauliflower mosaic virus (CaMV) 35S promoter, or more particularly the double enhanced cauliflower mosaic virus promoter, comprising two CaMV 35S promoters in tandem (referred to as a "Double 35S" promoter).

[0055] It may be desirable to use a tissue-specific or developmentally regulated promoter instead of a constitutive promoter in certain circumstances. A tissue-specific promoter allows for expression of immunoglobulins in certain tissues. Suitable fruit-specific promoters include the tomato E8 promoter (Deikman et al, (1988), EMBO J, 7, 3315-3320), 2A11 (Van Haaren et al, Plant Mol Biol, 21, 625-640), E4 (Cordes et al, (1989), Plant Cell, 1, 1025-1034) and PG (Bird et al, (1988), Plant Mol. Biol., 11, 651-662,) Nicholass et al, (1995), Plant Molecular Biology, 28, 423-435.

[0056] The invention provides in a further aspect an expression cassette comprising as operably linked components in the 5'-3' direction of transcription, a promoter functional in a plant cell, one or more DNA sequences encoding a heavy chain immunoglobulin or an active fragment or derivative thereof or one or more sequences encoding a protein functionally equivalent thereto and a transcriptional and translational termination regulatory region functional in a plant cell.

[0057] The promoter and termination regulatory regions will be functional in the host plant cell and may be heterologous (that is, not naturally occurring in the host plant) or homologous (derived from the plant host species) to the plant cell and the DNA sequence. Suitable promoters which may be used are described above.

[0058] The termination regulatory region may be derived from the 3' region of the gene from which the promoter was obtained or from another gene. Suitable termination regions which may be used are well known in the art and include *Agrobacterium tumefaciens nopaline synthase terminator (Tnos)*, *Agrobacterium tumefaciens mannopine synthase terminator (Tmas)* and the *CaMV 35S terminator (T35S)*. Particularly preferred termination regions for use according to the invention include the tobacco ribulose biphosphate carboxylase small subunit termination region (TrbcS) or the *Tnos* termination region.

[0059] Such DNA constructs may suitably be screened for activity by transformation into a host plant via *Agrobacterium* and screening for immunoglobulin levels.

[0060] In order to be able to select for plant cells that have integrated the construct, the expression construct may conveniently be joined to a marker which permits screening or selection, according to methods well known in the art.

[0061] Conveniently, the expression cassette according to the invention may be prepared by cloning the individual promoter/gene/ terminator units into a suitable cloning vector. Suitable cloning vectors are well known in the art, including such vectors as pUC (Norrander et al, (1983, Gene 26, 101-106), pEMBL (Dente et al (1983), Nucleic Acids Research, 11, 1645-1699), pBLUESCRIPT (available from Stratagene), pGEM (available from Promega) and pBR322 (Bolivar et al, (1977), Gene, 2, 95-113). Particularly useful cloning vectors are those based on the pUC series. The cloning vector allows the DNA to be amplified or manipulated, for example by joining sequences. The cloning sites are preferably in the form of a polylinker, that is a sequence containing multiple adjacent restriction sites, so as to allow flexibility in cloning.

[0062] Suitably, the nucleotide sequences for the genes may be extracted from the Genbank nucleotide database and searched for restriction enzymes that do not cut. These restriction sites may be added to the genes by conventional methods such as incorporating these sites in PCR primers or by sub-cloning.

[0063] Preferably the DNA construct according to the invention is comprised within a vector, most suitably an expression vector adapted for expression in an appropriate host (plant) cell. It will be appreciated that any vector which is capable of producing a recombinant plant comprising the introduced DNA sequence will be sufficient.

[0064] Suitable vectors are well known to those skilled in the art and are described in general technical references such as Pouwels et al, Cloning Vectors. A laboratory manual, Elsevier, Amsterdam (1986). Particularly suitable vectors include the Ti plasmid vectors.

[0065] Transformation techniques for introducing the DNA constructs according to the invention into host cells are well known in the art and include such methods as micro-injection, using polyethylene glycol, electroporation, or high velocity ballistic penetration. A preferred method for use according to the present invention relies on agrobacterium-mediated transformation.

[0066] After transformation of the plant cells or plant, those plant cells or plants into which the desired DNA has been incorporated may be selected by such methods as antibiotic resistance, herbicide resistance, tolerance to amino-acid analogues or using phenotypic markers.

[0067] Various assays, well known in the art, may be used to determine whether the plant cell shows the presence of the desired antibody, or fragment thereof. These include binding assay such as an ELISA or radio-immunoassay.

[0068] Plant cells transformed according to the invention may be grown in an appropriate culture or cultivated in soil or some other such suitable medium by methods well known in the art. It will be appreciated that the particular procedures adopted will vary depending on the plant species used, such variations being well within the knowledge of the average skilled practitioner.

[0069] These plant cells or plants may then conveniently be harvested and the desired antibody product processed using conventional extraction and purification techniques.

[0070] Seeds produced from modified plants containing the sequence coding for the desired antibody product can subsequently be grown to generate a progeny population of modified plants from which the desired product can be isolated.

[0071] Under some circumstances, it may be desirable to retain the antibody product with the plant rather than extracting and isolating the product. Where the modified plant or part thereof concerned is edible, for example, an antibody which has an effect on ingestion can conveniently be introduced in the form of a food product. In particular, edible plants or parts thereof, modified to incorporate an antibody or fragment or derivative thereof capable of binding a pre-selected antigen may conveniently be used in a method of passively immunising an animal, preferably a mammal and in particular a human, against said antigen. Particularly suitable antigens in this case will be pathogenic organisms or derivatives or parts thereof.

[0072] In addition to well-established application in the areas of diagnosis, therapy and purification, antibodies and fragments or derivatives thereof according to the present invention may suitably be used in methods to modify the properties of the plant in which they are produced. For example, the expression in a plant of functional anti-viral antibody can be useful in reducing pathogenicity and so provide a method of producing viral resistance. Alternatively, the introduction of antibodies into plants may be used to alter or interfere with plant metabolism by modulating the function of proteins present in the plant such as plant antigens and/or block phytohormones and metabolites.

[0073] The Invention will be further illustrated by means of the following examples which are provided by way of illustration only.

[0074] Techniques used for the modification and analysis of DNA materials were performed using standard procedures well known in the art, as described, for example, in Sambrook et al, "Molecular Cloning, A Laboratory Manual, Second Edition, Cold Spring Harbor, Laboratory Press 1989 (hereinafter Sambrook), unless otherwise indicated.

[0075] Restriction sites are underlined. HCV denotes heavy chain variable domain.

EXAMPLESExample 1: Construction of HCV expression vectors

[0076] The construction of the HCV plant expression plasmids involved several cloning steps. The anti-RR6 HCV gene sequence was isolated from male llamas immunised with BSA-RR6 (as described in WO 99/23221). The sequence of the DNA fragment used in the following example (HCV33: SEQ. ID. No. 1) is given in Figure 1.

pSJ.30

[0077] The plasmid pSJ.30 is a derivative of the binary vector pGPTV-KAN (Becker *et al.*, Plant Mol. Biol. 20: 1195-1197, 1992) modified as follows: An approximately 750 bp (Sac I, T4 DNA polymerase blunted - Sal I) fragment of pJIT60 (Guerineau *et al.*, Plant Mol. Biol. 18: 815-818, 1992) containing the duplicated cauliflower mosaic virus (CaMV) 35S promoter (denoted 2x35S; Cabb-JI strain, equivalent to nucleotides 7040 to 7376 duplicated upstream of 7040 to 7433, Frank *et al.*, Cell 21: 285-294, 1980) was cloned into the Hind III (klenow polymerase repaired) - Sal I sites of pGPTV-KAN to create pSJ.30.

pPV.3

[0078] The shuttle vector pPV.3 was constructed by removing the HindIII/EcoRI multiple cloning site from pUC19 and replacing it by a synthetic DNA fragment, destroying the original EcoRI and HindIII sites and introducing a SgfI, HindIII, KpnI, EcoRI and XbaI restriction site. The new insert was constructed by annealing the synthetic oligonucleotides PCR.624 (SEQ. ID. No. 12) and PCR.625 (SEQ. ID. No. 13) (Table 1) yielding the insert sequence:

PCR. 624

SgfI HindIII KpnI EcoRI XbaI

5' -AGCTGCGATCGCAAGCTTGGTACCGGGAATTCTCTAGA-3'

3' -CGCTAGCGTTTCGAACCATGGCCCTTAAGAGATCTTTAA-5'

PCR. 625

pPV.5

[0079] The KpnI/EcoRI insert from pSJ.30 containing the 2x35S-promoter sequence upstream of the GUS gene, followed by the Nos terminator sequence was inserted into the KpnI/EcoRI opened pPV.3 vector yielding the intermediate vector pPV.5.

pPV.5L

[0080] The GUS coding sequence was removed from the intermediate vector pPV.5 as a SalI/SacI fragment and replaced by a synthetic DNA fragment introducing a NcoI, NheI and MunI restriction site, while leaving the original SalI/SacI sites intact. The new insert was constructed by annealing the synthetic oligonucleotides PCR.626 (Seq. ID. No. 14) and PCR.627 (Seq. ID. No. 15) (Table 1) yielding the insert sequence:

PCR. 626

SalI NcoI NheI MunI SacI

5' -TCGACCCATGGCCCGCTAGCCAATTGGAGCT-3'

3' -GGGTACCGGGCGATCGGTAAACC-5'

PCR. 627

pPV.5LN

[0081] The sequence present immediately 5' of the start codon ATG in this vector (CCACCATGG) was replaced by the plant Kozak sequence TAAACCATGG using PCR. Oligonucleotides PCR.640 (Seq. ID. No. 16) and PCR.641 (Seq. ID. No. 17) (Table 1) was used to amplify the 189 bp 3' fragment of the 2x355 promoter from vector pCP031 (Engelen et. al., Plant Mol. Biol., 26, 1701-1710, 1994), modifying the Kozak sequence via PCR.641. pCP031 and the amplified fragment (SEQ. ID. No. 18) were digested with HindIII/EcoRV and EcoRV/NcoI respectively and used to replace the promoter present in the intermediate vector pPV.5L. After verification of the sequence by sequencing with primer Seq. ID. No. 19 the vector pPV.SLN was used in subsequent cloning steps.

PCR.640/PCR.641 amplification product (SEQ. ID. No. 18)

PCR.640

CCACCCACGAGGGAACATCGTGGAAAAAGAAGACGTTCCAACCACGTCTTCAAAGCAAGT
GGTGGGTGCTCCCTTGTAGCACCTTTTTCTTCTGCAAGGTTGGTGCAGAAAGTTTCGTTCA

EcoRV

GGATTGATGTGATATCTCCACTGACGTAAGGGATGACGCACAATCCCACTATCCTTCGCA
CCTAACTACACTATAGAGGTGACTGCATTCCCTACTGCTTGTTAGGGTGATAGGAAGCGT

NcoI

AGACCCTTCCTTTATATAAGGAAGTTCATTTCAATTTGGAGAGGACCTCGAGTGTAACCA

TCTGGGAAGGAAATATATTCCTTCAAGTAAAGTAAACCTCTCCTGGAGCTCACATTTGGT

PCR.641

TGGGAATTC**ACCCTTAAG**pPV.5LN-PRIa-HCV33-myc

[0082] The PstI/EcoRI HCV33-myc fragment from pPIC.HCV33-myc (described in WO 99/23221) was inserted into the NcoI/MunI opened vector pPV.5LN together with a 105 bp NcoI/PstI fragment encoding the PRIa leader sequence allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment.

[0083] The NcoI/PstI PRIa leader (Firek et. al., Plant Mol. Biol., 23, 861-870, 1993) sequence (including the first 4 amino acids of the HCV fragment (QVQL)) was constructed by annealing the synthetic oligonucleotides PCR.293 (Seq. ID. No. 20), PCR.294 (Seq. ID. No. 21), PCR.295 (Seq. ID. No. 22), PCR.296 (Seq. ID. No. 23), and PCR.297 (Seq. ID. No. 24) (Table 1). The product of the annealing reaction was cloned as a HindIII/PstI fragment into pUC.19. After verification of the sequence of the synthetic gene fragment by sequencing with primers Seq. ID. No. 25, 26 the NcoI/PstI fragment was excised and used in subsequent cloning reactions.

NcoI/PstI fragment encoding PRIA leader sequence (DNA Sequence Seq. ID. No. 27, Protein Sequence Seq. ID. No. 28)

[0084]

5

NcoI

10

CCATGGGATTTGTTCTCTTTTCACAATTGCCTTCATTTCTTCTGTCTCTACACTTCTCTT
GGTACCCTAAACAAGAGAAAAGTGTTAACGGAAGTAAAGAAGAACAGAGATGTGAAGAGAA
M G F V L F S Q L P S F L L V S T L L L

15

PstI

ATTCCTAGTAATATCCCACTCTTGCCGTGCCAGGTGCAGCTGCAG
TAAGGATCATTATAGGGTGAGAACGGCACGGGTCCACGTCGACGTC
F L V I S H S C R A Q V Q L

20

pPV.5LN-GBSS-HCV33-myc

25 **[0085]** The PstI/EcoRI HCV33-myc fragment from pPIC.HCV33-myc was inserted into the NcoI/MunI opened vector pPV.SLN together with a 252 bp NcoI/PstI fragment encoding the GBSS (including the first 4 amino acids of the HCV fragment (QVQL)) leader sequence allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment.

30 **[0086]** The NcoI/PstI GBSS leader sequence was constructed by amplifying potato tuber cDNA (Sp. Desiree) with synthetic oligonucleotides 132 (Seq. ID. No. 29) and 133 (Seq. ID. No. 30), 134 (Seq. ID. No. 31) and 136 (Seq. ID. No. 32) (Table 1). The products of the amplification reaction were cloned as a NcoI/HpaI, HpaI/PstI fragment into pGEM5Zf(+) (Promega). After verification of the sequence by sequencing with primers Seq. ID. No. 26, 33 (Table 1) the NcoI/PstI was excised and used in subsequent cloning reactions.

35

40

45

50

55

NcoI/PstI fragment encoding GBSS leader sequence (DNA Sequence Seq. ID. No. 34, Protein Sequence Seq. ID. No. 35)

[0087]

NcoI

CCATGGCAAGCATCACAGCTTCACACCACTTTGTGTCAAGAAGCCAACTTCACTAGACACCA
GGTACCGTTTCGTAGTGTCTGAAGTGTGGTGAAACACAGTTCTTCGGTTTGAAGTGATCTGTGGT
M A S I T A S H H F V S R S Q T S L D T

AATCAACCTTGTACAGATAGGACTCAGGAACCATACTCTGACTCACAATGGTTTAAGGGCT
TTAGTTGGAACAGTGTCTATCCTGAGTCCTTGGTATGAGACTGAGTGTTACCAAATCCCCGA
K S T L S Q I G L R N H T L T H N G L R A

GTTAACAACTTGATGGGCTCCAATCAAGAACTAATACTAAGGTAACACCCAAGATGGCATC
CAATTGTTTGAACACCCGAGGTTAGTTCTTGATTATGATTCCATTGTGGGTTCTACCGTAG
V N K L D G L Q S R T N T K V T P K M A S

PstI

CAGAACTGAGACCAAGAGACCTGGATGCTCAGCTACCATTGTTTGTGGAAAACAGGTGCAGCTGCAG
GTCTTGACTCTGGTTCTCTGGACCTACGAGTCGATGGTAACAAACACCTTTTGTCCACGTCGACGTC
R T E T K R P G C S A T I V C G K Q V Q L Q

pPV.5LN-HCV33-myc

[0088] The PstI/EcoRI HCV33-myc fragment from pPIC.HCV33-myc was inserted into the NcoI/MunI opened vector pPV.5LN together with a synthetic NcoI/PstI DNA fragment allowing the in-frame fusion with the ATG start codon. The synthetic linker was constructed by annealing the synthetic oligonucleotides PCR.652 (Seq. ID. No. 36) and PCR.653 (Seq. ID. No. 37) (Table 1) yielding the insert sequence:

PCR. 652

NcoI PstI
5' -CATGCAGGTGCAGCTGCA-3'
3' -GTCCACGTCG-5'

PCR. 653

pPV.5LN-HCV33-myc-SEKDEL

[0089] The HindIII/BstEII Promoter-leader-HCV33 fragment inserts from pPVSLN-PRla-HCV33-myc, pPVSLN-GBSS-HCV33-myc and pPV5LN-no leader-HCV33-myc was cloned into HindIII/MunI opened pPV5LN-PRla-HCV33-myc vector together with a myc-SEKDEL encoding BstEII/EcoRI PCR fragment thus introducing the sequence SEKDEL at the C-terminal end of the myc-tag, thus yielding pPV.5LN-PRla-HCV33-myc-SEKDEL, pPV.5LN-GBSS-HCV33-myc-SEKDEL and pPV.5LN-No leader-HCV33-myc-SEKDEL. The PCR fragment was generated using the synthetic oligo-

nucleotides PCR.300 (Seq. ID. No. 38) and PCR.690 (Seq. ID. No. 39) (Table 1) using pPV.5LN-PR1a-HCV33-myc as a template.

5 pSJ.34

[0090] The BamHI restriction site of the pGPTV-KAN expression vector containing the kanamycin resistance selectable marker gene (Becker *et. al.*, Plant Mol. Biol., 20, 1195-1197, 1992) was destroyed by restricting with BamHI followed by a Klenow reaction.

10 pPV.8 plant transformation vectors

[0091] The HindIII/EcoRI inserts from vectors pPV.5LN-PR1a-HCV33-myc, pPV.5LN-GBSS-HCV33-myc, pPV.5LN-No leader-HCV33-myc, pPV.5LN-PR1a-HCV33-myc-SEKDEL, pPV.5LN-GBSS-HCV33-myc-SEKDEL and pPV.5LN-No leader-HCV33-myc-SEKDEL were inserted into the HindIII/EcoRI vector fragment of pSJ.34, yielding pPV.8-GBSS-HCV33-myc, pPV.8-No leader-HCV33-myc, pPV.8-PR1a-HCV33-myc-SEKDEL, pPV.8-GBSS-HCV33-myc-SEKDEL and pPV.8-No leader-HCV33-myc-SEKDEL respectively.

pPV.5LN-2x35S-GBSS-scFv3299-hydrophilll

[0092] The PstI/EcoRI (partial) fragment containing scFv3299-hydrophilll (also known as scFv.KC in patent filing WO 96/27612 and described in more detail in WO 99/27386) was inserted into the NcoI/MunI opened vector pPV.5LN together with a 252 bp NcoI/PstI fragment encoding the GBSS leader sequence (as described for pPV.5LN-GBSS-HCV33-myc) allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment. By way of explanation, the sequence of scFv3299 is disclosed in W096/27612 (wherein it is described as FvKC-II), but with the BstEII-SacI linker fragment replaced with a BstEII-SacI fragment encoding a flexible linker yielding the sequence: TVTVSS-GGGGSGGGGSGGGGSDIELT (Seq. ID. No. 40). The DNA sequence of the linker is described by Jackson *et. al.* (Selection of variants of antibodies and other protein molecules using phage display on the surface of bacteriophage fd.: in Protein engineering: A Practical Approach, eds. Rees, Sternberg and Wetzel, publ. IRL Press, Oxford, 1992)

30 pSJ.34-2x35S-GBSS-scFv3299-hydrophilll

[0093] The HindIII/EcoRI 2x35S promoter-GBSS-scFv3299-hydrophilll fragment of pPV.5LN-2x35S-GBSS-scFv3299-hydrophilll was inserted into the HindIII/EcoRI vector fragment of pSJ.34.

35 pSJ.89

[0094] A HindIII/SacI fragment containing (1x) 35S promoter-GUS.int (Vancanneyt *et. al.*, Mol. Gen. Genet, 220, 245-250, 1990) was inserted into the HindIII/SacI vector fragment of pSJ.34. This plant transformation vector with Kan resistance contains the 35S promoter driving the GUS.int gene.

40 pSJ.103

[0095] The CERV promoter was isolated by PCR from infected *Dianthus barbatus* leaf material obtained from Dr Rene van der Lugt, Research Institute for Plant Protection (IPO-DLO), Binnehaven 5, P.O. Box 9060, NL-6700 GW, Wageningen, Netherlands, using the primers:

[0096] The approximately 380 bp fragment (corresponding to nucleotides 6737-7118 of Hull *et. al.*, EMBO Journal, 5, 3083-3090, 1986) was amplified with oligonucleotides CERV1 (Seq. ID. No. 41) and CERV2 (14 nucleotides downstream of the transcription start site) (Seq. ID. No. 42) (Table 1) and was cloned into the TA cloning vector pT7Blue (Invitrogen) with the promoter in the same orientation as the T7 promoter. Sequence analysis showed that the isolated CERV promoter differed from the published sequence at several positions (T-C at 6790, C-T at 6826, A-G at 6872 and T-A at 6729) the most significant of which was a small deletion in a polyA tract in the 5' untranslated leader (4A's vs. 9A's in the published sequence). These changes probably represent differences in virus isolates rather than PCR errors but these cannot be ruled out.

[0097] A HindIII - BamHI fragment from pSJ.103 was used as the source of the promoter in plant transformation vectors. Since the HindIII site was 13 nucleotides downstream of the CERV1 oligonucleotide, this sequence was not present in the promoter.

pSJ.108

[0098] The HindIII/BamHI CERV promoter fragment of pSJ.103 was inserted into the HindIII/BamHI vector fragment of pSJ.89. This plant transformation vector with Kan resistance contains the CERV promoter driving the GUS.int gene.

pPV.7

[0099] The HindIII/EcoRI insert from pSJ.108 containing the CERV promoter sequence upstream of the GUS gene, followed by the Nos terminator sequence was inserted into the HindIII/EcoRI opened pPV.3 vector yielding the intermediate vector pPV.7.

pPV.7L

[0100] The GUS coding sequence was removed from the intermediate vector pPV.7 as a BamHI/SacI fragment and replaced by a synthetic DNA fragment introducing a NcoI, NheI and MunI restriction site, while leaving the original BamHI/SacI sites intact. The new insert was constructed by annealing the synthetic oligonucleotides PCR.628 (Seq. ID. No. 43) and PCR.629 (Seq. ID. No. 44) (Table 1) yielding the insert sequence:

PCR. 628

BamHI NcoI NheI MunI SacI
 5' -GATCCCATGGCCCGCTAGCCAATTGGAGCT-3'
 3' -GGTACCGGGCGATCGGTTAACC-5'

PCR. 629

pPV.7LN

[0101] The sequence present immediately 5' of the start codon ATG in this vector (GATCCCATGG) was replaced by the plant Kozak sequence TAAACCATGG by replacing the BamHI/NheI fragment present in pPV.7L with a synthetic fragment constructed by annealing the synthetic oligonucleotides PCR.645 (Seq. ID. No. 45) and PCR.646 (Seq. ID. No. 46) (Table 1) yielding the vector pPV.7LN.

PCR. 645

BamHI NcoI NheI
 5' -GATCCACCTCGAGTGTAACCATGGCCCG-3'
 3' -GTGGAGCTCACATTTGGTACCGGGCGATC-5'

PCR. 646

pPV.7LN-GBSS-HCV33-myc

[0102] The PstI/MunI HCV33-myc fragment of pPIC.HCV33-myc was inserted into the NcoI/MunI opened vector pPV.7LN together with a 252 bp NcoI/PstI fragment encoding the GBSS leader sequence (as described for pPV.5LN-GBSS-HCV33-myc) allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment.

pPV.7LN-GBSS-Clone1-His6-myc

[0103] The Clone1-His6-myc sequence was amplified by PCR using oligonucleotides PCR.288 (Seq. ID. No. 47) and JW HCV2 (Seq. ID. No. 48) (Table 1) from pHEN.Clone1-His6-myc. The PstI (partial)/MunI fragment was inserted

into the NcoI/MunI opened vector pPV.7LN together with a 252 bp NcoI/PstI fragment encoding the GBSS leader sequence (as described for pPV.SLN-GBSS-HCV33-myc) allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment. The DNA sequence of clone 1 (Seq. ID. No 5, Figure 6) was then verified with sequencing primers, Seq. ID. No. 19,49 (Table 1).

pPV.7LN-GBSS-Clone46-His6-myc

[0104] The Clone46-His6-myc sequence was amplified by PCR using oligonucleotides PCR.288 and JW HCV2 from pHEN.Clone46-His6-myc. The PstI/MunI fragment was inserted into the NcoI/MunI opened vector pPV.7LN together with a 252 bp NcoI/PstI fragment encoding the GBSS leader sequence (as described for pPV.5LN-GBSS-HCV33-myc) allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment. The DNA sequence of clone 46 (Seq. ID. No. 6, Figure 12) was then verified by sequencing with primers, Seq. ID. No. 19,49 (Table 1).

pPV.7LN-Clone18-His6-myc

[0105] The Clone18-His6-myc sequence was amplified by PCR using oligonucleotides PCR.288 and JW HCV2 from pHEN.Clone18-His6-myc. The NcoI/MunI fragment was inserted into the NcoI/MunI opened vector pPV.7LN allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment. The DNA sequence of clone 18 (Seq. ID. No. 8, Figure 14) was then verified by sequencing with primers, Seq. ID. No. 19,49 (Table 1).

pPV.7LN-Clone18-His6-myc-SEKDEL

[0106] The Clone18-His6-myc sequence was amplified by PCR using oligonucleotides PCR.288 and JW HCV1 (Seq. ID. No. 50) (Table 1; to generate 3' SEKDEL endoplasmic retention signal) from pHEN.Clone18-His6-myc. The NcoI/MunI fragment was inserted into the NcoI/MunI opened vector pPV.7LN allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment. The DNA sequence of Clone18-His6-myc-SEKDEL (Seq. ID. No. 10, Figure 15) was then verified by sequencing with primers, Seq. ID. No. 19,49 (Table 1).

pSJ.35

[0107] The BamHI restriction site of the pGPTV-HYG expression vector containing the hygromycin resistance selectable marker gene (Becker *et. al.*, Plant Mol. Biol., 20, 1195-1197, 1992) was destroyed by restricting with BamHI followed by a Klenow reaction.

pSJ.35-CERV plant transformation vectors

[0108] The HindIII/EcoRI inserts from vectors pPV.7LN-GBSS-HCV33-myc, pPV.7LN-GBSS-Clone1-His6-myc, pPV.7LN-GBSS-Clone46-His6-myc, pPV.7LN-Clone18-His6-myc and pPV.7LN-Clone18-His6-myc-SEKDEL were inserted into HindIII/EcoRI vector fragment of pSJ.35, yielding pSJ.35-CERV-GBSS-HCV33-myc, pSJ.35-CERV-GBSS-Clone1-His6-myc, pSJ.35-CERV-GBSS-Clone46-His6-myc, pSJ.35-CERV-Clone18-His6-myc and pSJ.35-CERV-Clone18-His6-myc-SEKDEL respectively.

pCJ.102

[0109] Oligonucleotides 167 (Seq. ID. No. 51) and 168 (Seq. ID. No. 52) (Table 1) were used to amplify the 460bp 5' fragment of the GUS gene (from vector pSJ.34) and modifying the 5' end to create a NcoI site and thus modifying the second amino acid.

The PCR amplified fragment was digested with NcoI and EcoRV and inserted into HindIII/EcoRI vector fragment of pSJ.34 together with the HindIII/NcoI 2x35S promoter fragment of pPV.5LN and the EcoRV/EcoRI 3'GUS-Nos terminator fragment of pSJ.34, yielding the vector pCJ.102. The DNA sequence of the amplified fragment was verified by sequencing with primer Seq. ID. No. 53 (Table 1).

pSJ.35-Lat52-Clone18-His6-myc

[0110] The NcoI/EcoRI Clone18-His6-myc-Nos fragment of pPV.7LN-Clone18-His6-myc was inserted into the Sall/EcoRI vector fragment of pSJ.35 together with the Sall/NcoI Lat52 promoter fragment of pLAT52-7 (Twell *et. al.*, Development, 109, 705-13, 1990).

pSJ.35-Lat52-Clone18-His6-myc-SEKDEL

[0111] The NcoI/EcoRI Clone18-His6-myc-SEKDEL-Nos fragment of pPV.7LN-Clone18-His6-myc-SEKDEL was inserted into the Sall/EcoRI vector fragment of pSJ.35 together with the Sall/NcoI Lat52 promoter fragment of pLAT52-7 (Twell *et. al.*, Development, 109, 705-13, 1990).

pSJ.35-Lat52-HCV33-myc-SEKDEL

[0112] The PstI/SacI HCV33-myc-SEKDEL fragment of pPV.8-GBSS-HCV33-myc-SEKDEL was inserted into the Sall/SacI vector fragment of pSJ.35 with the Sall/PstI Lat52 promoter-5' antibody gene fragment of pSJ.35-Lat52-Clone18-His6-myc.

Example 2: Transformation and cultivation of *Nicotiana Tabacum*

[0113] Cells of the agrobacterium strain LBA4404 were transformed with the described plasmids using the freeze thaw method (Plant Molecular Biology Manual PMAN A3/7).

[0114] Seeds of *Nicotiana Tabacum* (var. Petit Havana SR1) were surface sterilised for 10 minutes in a solution of 10% sodium hypochlorate, rinsed 3 times in sterile distilled water and planted in Murashige & Skoog (MS) basal medium + 3% sucrose + 0.9% agar.

[0115] After 2 weeks growth seedlings were thinned to 2 per vessel; sterile plantlets were maintained by monthly shoot cuttings onto MS basal media +3% sucrose + 0.9% agar.

[0116] Discs were punched from leaves of these plants using a sterile cork borer, then incubated for 10 minutes in a culture of the agrobacterium strain LBA4404 containing the described plasmids which had been grown overnight in Lennox Broth (5g/l NaCl, Yeast Extract 10 g/l, 10 g/l Bacto Tryptone, 15g/l agar). The overnight culture was spun down for 10 minutes at 3000G and the growth media replaced by re-suspending the cells in MS Basal medium +3% sucrose.

[0117] Discs were removed from the culture and blotted dry on sterile filter paper before plating face down on a nurse culture plate of tobacco cells. This plate was prepared by adding 2ml of a cell suspension culture of *Nicotiana Bethiana* onto a petri dish containing 25ml of MS Salts, B5 vitamins (1mg/l Nicotinic acid, 1mg/l Pyridoxine, 10mg/l Thiamine, 100mg/l Inositol) +1mg/l 2-4 D, 0.2 mg/l BAP & 0.8% agar. The cells were swirled to cover the agar then covered with a sterile filter paper disc.

[0118] Infected leaf discs were incubated at 26°C in a light intensity of 2000 lux for 2 days, then removed from the nurse culture plates to selection media (MS Basal media +3% sucrose, 0.2mg/l IAA, 1mg/l BAP, 0.9% agar + cefotaxime 500mg/l and kanamycin 100 mg/l or hygromycin B 20mg/l). Plates were then incubated at 3000 lux, 26 C, 16 hours day/8 hour night and transferred every 2 weeks to fresh media (as previously). Shoots that appeared from the cut edges of the discs were removed to MS Basal media + 3% sucrose, cefotaxime 500mg/l and kanamycin 100mg/l or hygromycin B 20mg/l to root. Leaf samples were taken from rooted shoots for testing as described in Example 3, 4, 5, 7, 8 & 9.

Example 3: Extraction and detection of HCV material

Extraction of leaf tissue for analysis of anti-RR6 antibody (denoted HCV 33)

[0119] Leaf tissue was homogenised with Polytron probe in 2ml microcentrifuge tubes in the presence of 4ml extraction buffer (200mM Tris-HCl pH7.5, 5mM EDTA, 0.1% Tween 80, 1 mM PMSF {PhenylMethylSulfonylFluoride}) per 1gr leaf material. Solid material was removed from the homogenate by centrifugation (10 min 13,000g). The cleared supernatant was stored at -20°C. Total protein concentration was determined using the PIERCE BCA Protein Assay Reagent detection system.

Extraction of tomato fruit tissue for analysis of anti-RR6 antibody (denoted HCV33)

[0120] Fruit were dissected into peel, flesh and columella (centre of fruit), frozen and ground with a mortar and pestle in liquid nitrogen. Extracts were prepared as for leaf tissue above.

Extraction of leaf tissue for analysis of anti-potato

SBEII antibodies (denoted Clone1 and Clone46)

[0121] Leaf tissue was homogenised with Polytron probe in 2ml microcentrifuge tubes in the presence of 4 ml SBE

extraction buffer (100mM Tris-HCL pH7.5, 5mM EDTA, 2.5mM 1,4-dithiothreitol, 0.1% (w/v) sodium metabisulphite, 5% (v/v) glycerol, 2.5% (w/v) polyvinylpyrrolidone, 1mM PMSF, 1 tablet Complete (Roche) per 50ml) per 1gr fresh weight. Solid material was removed from the homogenate by centrifugation (10 min 13,000g). The cleared supernatant was stored at -20°C. Total protein concentration was determined using the BioRad Protein Assay Reagent detection system.

Extraction of leaf tissue for analysis of anti-GUS antibody (denoted Clone18)

[0122] As described for analysis of anti-RR6 antibody, except GUS extraction buffer (50mM phosphate pH 7.0, 10mM EDTA, 0.1% Triton X-100, 1 mM DTT).

Detection of RR6 binding activity

[0123] 96 well ELISA plates (Greiner) were coated overnight at 4°C with 100µl per well of BSA-RR6 conjugate (10µg/ml) in 0.05 M NaCO₃ buffer pH9.5. Following three washes with tap water, the plates were incubated for 1 hour at 37°C with 250µl per well of a blocking solution of 3% Marvel in TBST (20mM Tris-HCL pH7.5, 140mM NaCl, 0.1% Tween 20). Test plant extracts and appropriate standards (purified, *P.pastoris* produced HCV33-myc standard) in duplicate were added to wells in a total volume of 100µl TBST, and the plates incubated at 4°C overnight. Unbound antibody fragment was removed by 6 washes with tap water. 100µl of the primary monoclonal anti-myc antibody No. 4111 (1µg/mL) in 1% Marvel/TBST were added to each well. Following incubation at 37°C for 1 hour, unbound antibody was removed by 6 washes with tap water and 100 µL of the secondary anti-mouse antibody conjugated alkaline phosphatase (1ug/mL) was added to each well. Following incubation at 37°C for 1 hour, unbound antibody was removed by 6 washes with tap water, and alkaline phosphatase activity was detected by adding 100µl of 1mg/ml pNPP in 1M diethanolamine/1mM MgCl₂. Plates were incubated at room temperature for up to 5 hours before reading at 410nm on a Dynex, MCX plate reader.

Detection of SBEII binding activity

[0124] As for RR6 binding activity, except:

[0125] 100µl/well of 5µg/ml recombinant potato SBEII was bound to the plate (Jobling *et al.*, Plant Journal, 18, 163-171, 1999) in PBS overnight at 4°C

[0126] Purified and quantified clone1-myc from E.coli was used as the standard.

[0127] Blocking buffer: 1% BSA in TBST, all incubations at room temperature.

[0128] 100µl/well of 1:2,000 dilution of rabbit anti-llama IgG, as the primary detection antibody.

[0129] 100µl/well of 1ug/mL goat anti-rabbit antibody conjugated alkaline phosphatase, as the secondary detection antibody.

SDS-PAGE and Western blotting

[0130] Plant extracts were analysed on 10% bis-acrylamide gel using the Bio-Rad mini-Protein II system according to manufacturers instructions. 40µg samples and appropriate standards (purified *P.pastoris* produced HCV33-myc) were diluted in 50µl 2xSDS-PAGE loading buffer (20% Glycerol, 20mM Tris-HCL pH 8.0, 0.5% SDS, 0.005% bromophenol blue). The samples were mixed, boiled (5 minutes) and centrifuged briefly before loading onto the gel. The gels were run at 100V for 1 hour in 1x running buffer (25mM Tris, 200mM glycine, 1% SDS).

[0131] Proteins were transferred onto nylon membranes (Millipore) by semi dry blotting at 25V, 80mA for 45 minutes in semi-dry blotting buffer (50mM Tris, 40mM glycine, 0.04% SDS, 20% methanol). The membranes were then dried overnight at room temperature. The membranes were incubated in 3% Marvel, TBST for 10 minutes then in 1ug/mL mouse anti-myc antibody in 1% Marvel, TBST for 60 minutes at room temperature with constant agitation. Unbound antibody was removed by washing 6 times in TBST at room temperature with constant agitation. The blot was then incubated with the secondary antibody (rabbit anti-mouse IgG conjugated to alkaline phosphatase) in 1% Marvel, TBST for 60 minutes at room temperature with constant agitation. Unbound antibody was removed by washing 6 times in TBST at room temperature with constant agitation. Bound antibody was detected by incubating the filters in 10mL of 30mg/mL NBT and 15mg/ml BCIP. Washing the filters with water stopped the colour reaction.

EM studies

Step 1: Sample preparation:

[0132] Leaf samples were fixed using 1% Paraformaldehyde+ 0.05% glutaraldehyde in 0.05M sodium phosphate buffer pH6.8, for 4hrs at 4°C, and then washed in phosphate buffer overnight at 4°C. Samples were embedded by first dehydrating the samples through alcohol, 50, 70, 90% (15mins) and finally absolute alcohol (2x30mins). They were then placed in hydrophilic resin [6 parts LRGold resin +_4 parts GMA (low acid) +0.1% benzoin ethyl ether (UV activator). Resin was changed several times over period of three days. Finally, samples were embedded, using freshly prepared resin, in flat bottomed plastic moulds, allowing the resin to polymerise for 24hrs at room temperature, under nitrogen gas, using UV light (360nm).

Step 2: Staining:

[0133] Ultrathin leaf tissue sections were collected on collodion (2% in amyl acetate) coated nickel e.m. grids. Grids were floated on 20µl aliquots of TBS/BSA [20mM Tris/HCL buffer pH7.6 + 0.23M NaCl +1 bovine serum albumin] for 20mins at room temperature, following by incubation with 20µl aliquots of primary antibody (anti-myc or anti-hydrophil II, 1µg/ml) in TBS/BSA +0.05% Tween 20 for 18hrs at room temperature. Grids were washed thoroughly using TBS. The primary antibody was detected by incubating the grids with 20µl aliquots of secondary antibody/gold conjugate diluted in TBS/BSA (Goat a Mouse IgG; 5nm diluted 1:200, 10nm diluted 1:100 20nm diluted 1:50) for 60mins at room temperature. The grids were washed thoroughly, using TBS followed by washes with deionised water and dried. Some grids were silver enhanced, prior to poststaining with 2% aqueous Uranyl Acetate, followed by Lead Citrate. Photographs were taken using a either a Jeol 100CX Mk2 transmission electron microscope at 60,000 magnification (Figure 4) or a Jeol 1220 transmission electron microscope at 40,000 magnification (Figure 11).

Assay of SBE activity

[0134] All protein samples were diluted to a concentration of 1mg/mL using SBE extraction buffer. SBE was assayed using the phosphorylase a stimulation assay (Hawker *et al.*, Arch. Biochem. Biophys., 160, 530-551, 1974). The reaction was carried out at 30°C for 1h on 50µg protein in a final volume of 200µl and contained 100mM 2-(N-morpholino) ethanesulphonic acid (MES), 50mM [U-14C]glucose-1-phosphate (370MBq mole⁻¹, ICN), 1.3U rabbit muscle phosphorylase a (Sigma), 50µl plant extract at 1mg/ml protein. Glucose polymers were precipitated and washed as described by Hawker *et al.*, Arch. Biochem. Biophys., 160, 530-551 (1974). The final pellet was dissolved in DMSO and counted with ReadySafe-scintillation cocktail (Beckmann).

Assay of GUS activity

[0135] All protein samples were diluted to a concentration of 1 mg/ml using GUS extraction buffer. 10µl samples were then added to 190µl of 1 x reaction buffer (50 mM phosphate pH 7.0, 0.1% Triton X-100, 1 mM DTT, 1 mM p-nitrophenyl β-D-glucuronide) contained in the well of a flat-bottomed microtitre plate. The reaction was left to proceed at 37°C until the wells turned visibly yellow. At this point, 80µl of stop solution (2.5M 2-amino-2-methyl propanediol) was added and the plate read at 410nm using a Dynex plate reader. GUS activity was calculated based on positive control wells to which known quantities of GUS were added.

Histochemical staining for GUS activity

[0136] Tobacco pollen was stained for GUS activity in microtitre wells for up to 1 hour in 1 mM 5-bromo-4-chloro-3-indolyl β-D-glucuronic acid according to the method of Jefferson *et al.*, EMBO Journal 6: 3901-7, (1987). The buffer was modified by the use of histochemical staining buffer comprising 100 mM sodium phosphate pH 7.0, 10 mM EDTA, and 0.1% (v/v) Triton X-100 (Stomp, In: *Editorial comments*, Vol. 16, No.5. Cleveland: united State Biochemical, 1990). To minimise spread of the stain to surrounding pollen grains, 5mM potassium ferricyanide was added to localise staining (Lojda, Histochemie 22: 347-61, 1970; Mascarenhas and Hamilton, Plant Journal 2: 405-8, 1992).

Example 4: Expression of HCV antigen binding activity in the cytoplasm

[0137] Small samples of the leaves of 10 individual Tobacco plants transformed with the expression vector pPV. 8-HCV33-myc were tested for the presence of anti-RR6 binding activity using ELISA. One of the tested samples showed low levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity present in this clone (No. 943.37)

using SDS-PAGE/Western blot analysis, ELISA and total protein concentration measurement showed that HCV33-myc was expressed as 0.001-0.002% of total protein present (Figure 2 and Table 2).

Example 5: Expression of HCV antigen binding activity in plastids

[0138] Small samples of the leaves of 15 individual Tobacco plants transformed with the expression vector pPV.8-GBSS-HCV33-myc were tested for the presence of anti-RR6 binding activity using ELISA. Seven of the tested samples showed high levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity present in these clones using SDS-PAGE/Western blot analysis, ELISA and total protein concentration measurement showed that HCV33-myc was expressed as 0.5-1.0% of total soluble protein present (Figure 3 and Table 2). From the Western blot it can be concluded that the construct is processed correctly and that the GBSS leader is removed from the HCV fragment. Localisation of the HCV33 material in the chloroplasts was confirmed via immuno-histology studies (Figure 4).

[0139] The fruit of 10 individual tomato plants transformed with the expression vector pPV.8-GBSS-HCV33-myc were picked 15 days post-breaker (red stage), dissected and tested for the presence of anti-RR6 binding activity using ELISA. 5 of tested fruit expressed levels of RR6-binding activity which, by comparison to a standard curve of purified HCV33 binding to RR6, were over 0.1% (one was over 0.2%) of total soluble protein (Figure 5 and Table 2).

[0140] Similar high levels of expression were obtained with an HCV antibody of different specificity. Small samples of the leaves of 30 individual tobacco plants transformed with the expression vector pSJ.35-CERV-GBSS-Clone1-His6-myc were tested for the presence of anti-SBEII binding activity using ELISA. 3 of the tested plants expressed levels of SBEII-binding activity over 0.1% total soluble protein (Figure 7 and Table 2).

Example 6: Disruption of plastid by scFv expression

[0141] Tobacco plants were transformed with expression vector pSJ34-2x35S-GBSS-scFv3299-hydrophilinIII. After initial detection of high expression of hCG binding activity, these levels were lost - expression was unstable (not shown). Furthermore, immuno-histology studies (Figure 8) suggested that scFv expression had an adverse affect on plastid structure and function, perhaps due to the aggregation of multiple scFv molecules at the plastid membrane.

Example 7: Expression of HCV antigen binding activity in the apoplastic space

[0142] Small samples of the leaves of 21 individual Tobacco plants transformed with the expression vector pPV.8-PR1a-HCV33-myc were tested for the presence of anti-RR6 binding activity using ELISA. 12 of the tested samples showed low levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity present using showed that HCV33-myc was expressed as 0.0002-0.00074% of total protein present (Figure 9 and Table 2).

Example 8: Expression of HCV33 containing the ER retention signal SEKDEL

[0143] Small samples of the leaves of 22, 23 and 12 individual Tobacco plants transformed with the expression vectors pPV.8-no leader-HCV33-myc-SEKDEL, pPV.8-PR1a-HCV33-myc-SEKDEL, pPV.8-GBSS-HCV33-myc-SEKDEL respectively were tested for the presence of anti-RR6 binding activity using ELISA. None, 16 and 3 of the tested samples respectively, showed medium to high levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity present in these clones using SDS-PAGE/Western blot analysis, ELISA and total protein concentration measurement showed that HCV33-myc was expressed as 0, 0.4 and 0.05% of total protein present respectively (Figure 10 and Table 2). Localisation of the HCV33 material, which was directed to the secretory pathway by using the PR1a leader sequence, in the ER was confirmed using immuno-histology studies (Figure 11).

Example 9: Effect of neutralising HCV expression on SBE activity in tobacco plants

[0144] Samples of the leaves of 30 individual tobacco plants transformed with expression vector pSJ.35-CERV-GBSS-Clone46-His6-myc were assayed for SBE activity. Compared to the effect of the HCV33 anti-RR6 antibody in 30 control plants transformed with expression vector pSJ.35-CERV-GBSS-HCV33-myc, the specific anti-potato SBEII antibody Clone46 significantly reduced the SBE activity in the tobacco plants (Figure 13), indicating that the antibody has neutralised the enzyme activity.

Example 10: Effect of neutralising HCV expression on GUS activity in tobacco leaves

[0145] A tobacco line already transformed with expression vector pCJ.102 and shown to contain GUS activity in leaf tissue was retransformed with the expression vector pSJ35-CERV-Clone18-His6-myc-SEKDEL. Samples from 27 in-

dividual transformed plants were assayed for GUS activity in comparison to control pCJ.102 tobacco lines that had not been retransformed. A graphical representation of the results from measuring GUS activity in control lines and in lines containing Clone 18 is shown in Figure 16. A difference between the two sets of data is clearly evident, indicating that the presence of the anti-GUS antibody has decreased GUS activity.

Example 11: Effect of neutralising HCV expression on GUS activity in tobacco pollen

[0146] The tobacco line *lat52-GUS*, containing the pollen specific promoter *lat52* driving *GUS* in a single copy homozygous state, was obtained as a gift from Dr. David Twell (Botany Department, University of Leicester, UK). This line was retransformed with expression vector pSJ.35-Lat52-Clone18-His6-myc-SEKDEL. Control lines were obtained by transforming *lat52-GUS* tobacco with pSJ.35-Lat52-HCV33-myc-SEKDEL. Pollen from 20 lines of each set of transformed tobacco were tested by histochemical staining for GUS activity. Scanned images showing the staining pattern obtained in a control line and a line containing anti-GUS VHH clone 18 are shown in Figure 17. It can quite clearly be seen that GUS activity has been 'knocked-out' by the presence of anti-GUS antibody. Half the pollen retains GUS activity (shown by blue staining) due to segregation of the transgene at meiosis.

Example 12: Expression of whole heavy chain immunoglobulins in plants

Construction of HCV33-IgG2b expression vectors

pPV.5LN-HCV33-IgG2b

[0147] A cDNA clone IgG2b (Vu *et. al.*, Mol. Imm., 34, 16-17, 1997) was amplified by PCR using oligonucleotides 304 (Seq. ID. No. 58) and 305 (Seq. ID. No. 59), 306 (Seq. ID. No. 60) and 310 (Seq. ID. NO. 61) (Table 1). The amplified fragments were digested with BstEII/HincII and HincII/EcoRI respectively and inserted into NcoI/MunI opened vector pPV.5LN together with the NcoI/BstEII HCV33 fragment from vector pPV.5LN-HCV33-myc, yielding the vector pPV.5-HCV33-IgG2b. The DNA sequence of the amplified fragment was verified by sequencing with primers Seq. ID. No. 19, 53, 58, 60 and 62 (Table 1).

[0148] The predicted amino acid sequence contained two mutations at amino acid numbers 167 and 369 (Figure 18) as compared to the published sequence by Woolven *et. al.*, Immunogen., 50, 98-101, 1999. Oligonucleotides 314 (Seq. ID. No. 63) and 307 (Seq. ID. No. 64) (Table 1) were designed to change the mutations at amino acids 167 (proline) and 369 (Stop codon) to a serine and lysine respectively.

[0149] The HCV33-IgG2b clone was then amplified by PCR using oligonucleotides 314 and 307 (Table 1). The amplified fragment was digested with SacI/EcoRI and inserted into HindIII/MunI opened vector pPV.5LN together with the HindIII/SacI fragment from vector pPV.5-HCV33-IgG2b, yielding the vector pPV.5LN-HCV33-IgG2b. The DNA sequence of the amplified fragment was verified by sequencing with primers Seq. ID. No. No. 19, 53, 58, 60 and 62 (Table 1) (Figure 18).

pPV.5LN-PR1a-HCV33-IgG2b

[0150] The BstEII/EcoRI HCV33-IgG2b fragment from vector pPV.5LN-HCV33-IgG2b was inserted into HindIII/EcoRI opened pPV.5LN together with the HindIII/BstEII promoter-PR1a-HCV33 fragment from vector pPV.5LN-PR1a-HCV33-myc.

pPV.5LN-GBSS-HCV33-IgG2b

[0151] The BstEII/EcoRI HCV33-IgG2b fragment from vector pPV.5LN-HCV33-IgG2b was inserted into HindIII/EcoRI opened pPV.5LN together with the HindIII/BstEII promoter-GBSS-HCV33 fragment from vector pPV.5LN-GBSS-HCV33-myc.

pPV.5LN-PR1a-HCV33-IgG2b-SEKDEL

[0152] Vector pPV.5LN-HCV33-IgG2b was amplified by PCR using oligonucleotides 306 and 308 (Seq. ID. No. 65) (Table 1; to generate 3' SEKDEL endoplasmic retention signal). The amplified fragment was digested with HincII/EcoRI and inserted into HindIII/EcoRI opened pPV.5LN together with the HindIII/HincII fragment from vector pPV.5LN-PR1a-HCV33-IgG2b. The DNA sequence of the amplified fragment was the verified by sequencing with primers Seq. ID. No. 19, 63 (Table 1) (Figure 19).

pPV.8 2x35S plant transformation vectors

[0153] The HindIII/EcoRI inserts from vectors pPV.5LN-HCV33-IgG2b, pPV.5LN-PR1a-HCV33-IgG2b, pPV.5LN-GB-SS-HCV33-IgG2b and pPV.5LN-PR1a-HCV33-IgG2b-SEKDEL were inserted into HindIII/EcoRI vector fragment of pSJ.34, yielding pPV.8-HCV33-IgG2b, pPV.8-PR1a-HCV33-IgG2b, pPV.8-GBSS-HCV33-IgG2b and pPV.8-PR1a-HCV33-IgG2b-SEKDEL respectively.

Detection of HCV33-IgG2b binding activity

[0154] As for RR6 binding activity, except:

[0155] Purified and quantified HCV33-IgG2b from tobacco plants was used as the standard.

[0156] 100µl/well of 1:2,000 dilution of rabbit anti-llama IgG was used as the primary detection antibody.

[0157] 100µl/well of 1ug/mL goat anti-rabbit antibody conjugated alkaline phosphatase was used as the secondary detection antibody.

Expression of HCV33-IgG2b antigen binding activity in the cytoplasm

[0158] Small samples of the leaves of 20 individual Tobacco plants transformed with the expression vector pPV.8-HCV33-IgG2b were tested for the presence of anti-RR6 binding activity using ELISA. 10 of the tested samples showed low levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed as 0.011-0.031% of total soluble protein present (Figure 20).

[0159] Three plants chosen at random were selfed and allowed to set seed, the next generation was germinated and after 6 weeks small samples of leaf, stem and root were tested for the presence of anti-RR6 binding activity using ELISA. Two of the tested plants showed increased levels of RR6 binding activity as compared to the original analysis. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed at its highest in stem tissues at a maximum level of 1.07% total protein present (Figure 21).

Expression of HCV33-IgG2b antigen binding activity in the plastids

[0160] Small samples of the leaves of 20 individual Tobacco plants transformed with the expression vector pPV.8-GBSS-HCV33-IgG2b were tested for the presence of anti-RR6 binding activity using ELISA. 10 of the tested samples showed low levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed as 0.034-0.050% of total soluble protein present (Figure 22).

[0161] Three plants chosen at random were selfed and allowed to set seed, the next generation was germinated and after 6 weeks small samples of leaf, stem and root were tested for the presence of anti-RR6 binding activity using ELISA. All of the tested samples showed increased levels of RR6 binding activity as compared to the original analysis. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed at its highest in leaf tissues at a maximum level of 9.26% total soluble protein present (Figure 23).

Expression of HCV33-IgG2b antigen binding activity in the apoplastic space

[0162] Small samples of the leaves of 20 individual Tobacco plants transformed with the expression vector pPV.8-PR1a-HCV33-IgG2b were tested for the presence of anti-RR6 binding activity using ELISA. 17 of the tested samples showed significant levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed as 0.021-0.157% of total soluble protein present (Figure 24).

[0163] Three plants chosen at random were selfed and allowed to set seed, the next generation was germinated and after 6 weeks small samples of leaf, stem and root were tested for the presence of anti-RR6 binding activity using ELISA. All of the tested samples showed increased levels of RR6 binding activity as compared to the original analysis. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed at very high levels in all tissues tested, with the highest level found in leaf at a maximum of 13% total soluble protein present (Figure 25).

Expression of HCV33-IgG2b antigen binding activity in the endoplasmic reticulum

[0164] Small samples of the leaves of 20 individual Tobacco plants transformed with the expression vector pPV.8-HCV33-IgG2b were tested for the presence of anti-RR6 binding activity using ELISA. 13 of the tested samples showed significant levels of RR6 binding activity. Detailed analysis of the anti-RR6 binding activity present using showed that HCV33-IgG2b was expressed as 0.02-0.165% of total soluble protein present (Figure 26).

[0165] Three plants chosen at random were selfed and allowed to set seed, the next generation was germinated

and after 6 weeks small samples of leaf, stem and root were tested for the presence of anti-RR6 binding activity using ELISA. All of the tested samples showed increased levels of RR6 binding activity as compared to the original analysis. Detailed analysis of the anti-RR6 binding activity showed that HCV33-IgG2b was expressed at very high levels in all tissues tested, with the highest level found in leaf at a maximum of 16.3% of total soluble protein present (Figure 27).

Example 13: Expression of anti-SBEII (clone68) antigen binding activity in tobacco directed to the plastids

Construction of CERV-GBSS-Clone68-myc expression vectors

pPV.7LN-GBSS-Clone68-His6-myc

[0166] The Clone68-His6-myc sequence was amplified by PCR using oligonucleotides PCR.288 (Seq. ID. No. 47) and JWHCV2 (Seq. ID. No. 48) (Table 1) from pHEN.Clone68-His6-myc. The PstI/MunI fragment was inserted into the NcoI/MunI opened vector pPV.7LN together with a 252 bp NcoI/PstI fragment encoding the GBSS leader sequence (as described for pPV.5LN-GBSS-HCV33-myc) allowing the in-frame fusion with the 5' framework-1 sequences of the HCV fragment to create pPV.7LN-GBSS-Clone68-His6-myc. The DNA sequence of clone 68 was then verified with sequencing primers, Seq. ID. No. 19,49 (Table 1) (Figure 28).

pSJ.35-CERV plant transformation vector

[0167] The HindIII/EcoRI inserts from vector pPV.7LN-GBSS-clone68-His6-myc were inserted into HindIII/EcoRI vector fragment of pSJ.35, yielding vector pSJ.35-CERV-GBSS-clone68-His6-myc.

Detection of SBEII binding activity

[0168] This was performed as described in example 3 for anti-SBEII binding activity, except that purified and quantified clone68-His6-myc from *E.coli* was used as the standard.

Expression of anti-SBEII antigen binding activity in the plastids

[0169] High levels of expression were obtained with another HCV antibody with anti-SBEII antigen binding specificity. Small samples of the leaves of 30 individual tobacco plants transformed with the expression vector pSJ.35-CERV-GBSS-Clone68-His6-myc were tested for the presence of anti-SBEII binding activity using ELISA. 26 of the tested plants expressed levels of SBEII-binding activity above the standards with expression levels at a minimum of 0.3% total soluble protein (Figure 29).

Table 1: DNA sequence of the synthetic oligonucleotides used in the examples.

Code	SEQ. ID. No.	Nucleotide Sequence
PCR. 624	12	AGCTGCGATCGCAAGCTTGGTACCGGGAATTCTCTAGA
PCR. 625	13	AATTCTAGAGAATTCCCGGTACCAAGCTTGCTTGCATCGC
PCR. 626	14	TCGACCCATGGCCCGCTAGCCAATTGGAGCT
PCR. 627	15	CCAATTGGCTAGCGGGCCATGGG
PCR. 640	16	CCACCCACGAGGGAACATCGTG
PCR. 641	17	GAATTCCTCATGGTTTACACTCGAGGTCCTCTCCAAATGA
172	19	CCGGCAACAGGATTCAATCTT
PCR. 293	20	AGCTCCATGGGATTTGTTCTCTTTTACAATTGCCTTCAT
PCR. 294	21	TTCTTCTTGTTCTCTACACTTCTCTTATTCTAGTA
PCR. 295	22	ATATCCCACTCTTGCCGTGCCAGGTGCAGCTGCA
PCR. 296	23	GCTGCACCTGGGCACGGCAAGAGTGGGATATTACTAGGAATAAGAGAA
PCR. 297	24	GTGTAGAGACAAGAAGAAATGAAGGCAATTGTGAAAAGAGAACAAATCCC ATGG
152	25	GGAAACAGCTATGACCATGATTAC
191	26	TTTCCCAGTCACGACGTTGT
132	29	GACATCCCATGGCAAGCATCA
133	30	AAGCTTGTTAACAGCCCTTAA
134	31	AGGGCTGTTAACAACTTGAT
136	32	AGACTGCTGCAGCTGCACCTGCTTTCCACAAACAATGGTAGCTG
197	33	AATTAACCCCTCACTAAAGG
PCR. 652	36	CATGCAGGTGCAGCTGCA
PCR. 653	37	GCTGCACCTG
PCR. 300	38	CGCAAGACCCTTCCTCTATATAAG
PCR. 690	39	GAGCTCGAATTCTTATTATAGCTCATCTTTCTCTGAATTCAGATCCTCTT CTGAGATGAG
CERV1	41	ATCCTCAACTTCCAATCAGA
CERV2	42	TTCTTGAGAGATAGCTTGA
PCR. 628	43	GATCCCATGGCCCGCTAGCCAATTGGAGCT
PCR. 629	44	CCAATTGGCTAGCGGGCCATGG

PCR. 645	45	GATCCACCTCGAGTGTAACCATGGCCCG
PCR. 646	46	CTAGCGGGCCATGGTTTACACTCGAGGTG
PCR. 288	47	ATTGCCTACGGCAGCCGCTG
JWHCV2	48	TCCAACCAATTGTTACTATGCGGCCCCATTAGATCCTCTTCTGAG ATGAG
164	49	GTCTGTCTAAAGTAAAGTAGATGCG
JWHCV1	50	TCCAACCAATTGTTATCATAGCTCATCTTTCTCACTATTCAGATCC TCTTCTGAGATGAG
167	51	AGTCCCCCATGGTACGTCCTGTAGAAACC
168	52	CGTTTTTCGTCGGTAATCACCATTCC
161	53	CGCAAGACCCTTCCTTTATATAAG
304	58	AGGGGACCCAGGTCACCGTCTCCTCAGAACC
305	59	GAGCTTTGTTGTTGACCTTGCATTGAACTCTTTCCC
306	60	CAAATGCAAGGTCAACAACAAGCTC
310	61	TTCGAGCTCGAATTCTTATTATTTACCCGAAGACTGGGTGAT
303	62	CTGAGGAGACGGTGACCTGGGTCCCCT
314	63	AGCCCCTGAGCTCCTGGGAGGGCCCTCAGTCTTCATCTTCCCCCG
307	64	TTCGAGCTCGAATTCTTATTATTTACCCGAAGACTGGGTGATGGATTCT GGGTGTAGTGG
308	65	TTCGAGCTCGAATTCTTATTATAGCTCATCTTTCTCAGATTACCCGAAG ACTGGGTGATGGATTCTGGGTGTAGTGG

Table 2:

Summary of HCV expression levels in tobacco (unless otherwise stated) detected. Maximum observed expression levels are expressed as percentage of active antibody in the total amount of soluble protein extracted.

Construct	Compartment	Expression Level
pPV.8-no leader-HCV33-myc	Cytosol	0.002%
pPV.8-GBSS-HCV33-myc	Chloroplast	1.0%
pPV.8-PR1a-HCV33-myc	Apoplast	0.00074%
pPV.8-no leader-HCV33-myc-KDEL	Cytosol	<0.001%
pPV.8-GBSS-HCV33-myc-KDEL	Chloroplast	0.05%
pPV.8-PR1a-HCV33-myc-KDEL	ER	0.4%
pPV.8-GBSS-HCV33-myc	Plastid (Tomato fruit columella)	0.2%
pSJ.35-CERV-GBSS-Clonel-His6-myc	Chloroplast	0.2%

SEQUENCE LISTING

5 <110> Unilever plc
Unilever NV

10 <120> Production of Antibodies

<130> T7060

15 <160> 67

20 <170> PatentIn version 3.0

<210> 1

25 <211> 440
<212> DNA
<213> Artificial

30 <220>
<223> VHH with peptide linker
<220>

35 <221> CDS
<222> (1)..(417)

40 <400> 1
cag gtg cag ctg cag gag tca ggg gga gga ttg gtg cag gct ggg ggc 48
Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
tct ctg aga ctc tcc tgt gca gcc tgc gga cgc gcc acc agt ggt cat 96
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His
20 25 30
ggt cac tat ggt atg ggc tgg ttc cgc cag gtt cca ggg aag gag cgt 144
Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg
35 40 45
gag ttt gtc gca gct att agg tgg agt ggt aaa gag aca tgg tat aaa 192
Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys
50 55 60
gac tcc gtg aag ggc cga ttc acc atc tcc aga gat aac gcc aag act 240
Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr
65 70 75 80

EP 1 118 669 A2

	acg gtt tat ctg caa atg aac agc ctg aaa cct gaa gat acg gcc gtt	288
	Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val	
	85 90 95	
5	tat tat tgt gcc gct cga ccg gtc cgc gtg gat gat att tcc ctg ccg	336
	Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro	
	100 105 110	
	gtt ggg ttt gac tac tgg ggc cag ggg acc cag gtc acc gtc tcc tca	384
10	Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser	
	115 120 125	
	gaa caa aaa ctc atc tca gaa gag gat ctg aat taataagggc taagctcgaa	437
	Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn	
	130 135	
15	ttc	440
	<210> 2	
	<211> 139	
20	<212> PRT	
	<213> Artificial	
25	<400> 2	
	Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly	
	1 5 10 15	
30	Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His	
	20 25 30	
	Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg	
35	35 40 45	
	Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys	
	50 55 60	
40	Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr	
	65 70 75 80	
	Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val	
45	85 90 95	
	Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro	
	100 105 110	
50	Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser	
	115 120 125	
	Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn	
55	130 135	

EP 1 118 669 A2

```

<210> 3
<211> 11
5 <212> PRT
<213> Artificial

10 <220>
<223> myc linker
<400> 3
15 Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn
1 5 10

<210> 4
<211> 471
20 <212> DNA
<213> Artificial

25 <220>
<223> VHH with linker
<220>
30 <221> CDS
<222> (1)..(459)

35 <400> 4
cag gtg cag ctg cag cag tca ggg gga ggc ttg gtg cag gct ggg ggg 48
Gln Val Gln Leu Gln Gln Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15

tct ctg aga ctc tcc tgt gta gct tct gaa agc agc ttc agc aac aat 96
Ser Leu Arg Leu Ser Cys Val Ala Ser Glu Ser Ser Phe Ser Asn Asn
20 25 30

cac atg ggc tgg tac cgc cgg gct cca ggg aac cag cgc gag ctg gtc 144
His Met Gly Trp Tyr Arg Arg Ala Pro Gly Asn Gln Arg Glu Leu Val
35 40 45

gca act att agt cct ggt ggt agc aca cac tat gta gac tcc gtg aag 192
Ala Thr Ile Ser Pro Gly Gly Ser Thr His Tyr Val Asp Ser Val Lys
50 55 60

ggc cga ttc acc atc tcc cga gac aac gcc aag aac aca gtg tat cta 240
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr Leu
65 70 75 80

caa atg gac agc ctg aaa cca gag gac acg gcc gtc tat tac tgt gct 288
Gln Met Asp Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

55 gcc aag ggg agg ggg ctg cag gct atg cag tac tgg gcc cag ggg acc 336
Ala Lys Gly Arg Gly Leu Gln Ala Met Gln Tyr Trp Gly Gln Gly Thr

```

EP 1 118 669 A2

	100	105	110	
	ctg gtc acc gtc tcc tca gcg cac cac agc gaa gac ccc agc tcc gcg			384
	Leu Val Thr Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala			
5	115	120	125	
	gcc gcc cat cac cat cac cat cac ggg gcc gca gaa caa aaa ctc atc			432
	Ala Ala His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile			
	130	135	140	
10	tca gaa gag gat ctg aat ggg gcc gca tagtaacaat tg			471
	Ser Glu Glu Asp Leu Asn Gly Ala Ala			
	145	150		
	<210> 5			
15	<211> 153			
	<212> PRT			
	<213> Artificial			
20				
	<400> 5			
25	Gln Val Gln Leu Gln Gln Ser Gly Gly Gly Leu Val Gln Ala Gly Gly			
	1	5	10	15
	Ser Leu Arg Leu Ser Cys Val Ala Ser Glu Ser Ser Phe Ser Asn Asn			
	20	25	30	
30	His Met Gly Trp Tyr Arg Arg Ala Pro Gly Asn Gln Arg Glu Leu Val			
	35	40	45	
	Ala Thr Ile Ser Pro Gly Gly Ser Thr His Tyr Val Asp Ser Val Lys			
35	50	55	60	
	Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr Leu			
	65	70	75	80
40	Gln Met Asp Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala			
	85	90	95	
	Ala Lys Gly Arg Gly Leu Gln Ala Met Gln Tyr Trp Gly Gln Gly Thr			
45	100	105	110	
	Leu Val Thr Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala			
	115	120	125	
50	Ala Ala His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile			
	130	135	140	
	Ser Glu Glu Asp Leu Asn Gly Ala Ala			
55	145	150		

EP 1 118 669 A2

<210> 6
 <211> 468
 5 <212> DNA
 <213> Artificial

 10 <220>
 <223> VHH with linker
 <220>
 15 <221> CDS
 <222> (1)..(456)

 20 <400> 6
 cag gtg cag ctg cag gag tct ggg gga gcc ctg gtg cag gct ggg ggg 48
 Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
 1 5 10 15
 tct ctg aga ctc tcc tgt gta gcc tct gga aac acc ttc agt atc ata 96
 Ser Leu Arg Leu Ser Cys Val Ala Ser Gly Asn Thr Phe Ser Ile Ile
 25 20 25 30
 gct atg gcc tgg tac cgc cag gct cca ggg aag cag cgc gag gtg gtc 144
 Ala Met Ala Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Val Val
 35 35 40 45
 gca agt att aat agt att ggc agc aca aat tat gca gac tcc gtg aag 192
 Ala Ser Ile Asn Ser Ile Gln Ser Thr Asn Tyr Ala Asp Ser Val Lys
 50 55 60
 ggg cga ttc acc atc tcc aga gac aac gcc aag aac aca gtg tat ctg 240
 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr Leu
 35 65 70 75 80
 caa atg agc agc ctg aaa cct gag gac acg gcc gtc tat tac tgt gct 288
 Gln Met Ser Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 gcc ggt aat ttg ctg gtt aag agg cct tac tgg gcc cag ggg acc ctg 336
 Ala Gly Asn Leu Leu Val Lys Arg Pro Tyr Trp Gly Gln Gly Thr Leu
 40 100 105 110
 gtc acc gtc tcc tca gaa ccc aag aca cca aaa cca caa cca gcg gcc 384
 Val Thr Val Ser Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Ala Ala
 45 115 120 125
 gcc cat cac cat cac cat cac ggg gcc gca gaa caa aaa ctc atc tca 432
 Ala His His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser
 130 135 140
 50 gaa gag gat ctg aat ggg gcc gca tagtaacaat tg 468
 Glu Glu Asp Leu Asn Gly Ala Ala
 145 150

 55 <210> 7
 <211> 152

EP 1 118 669 A2

<212> PRT

<213> Artificial

5

<400> 7

Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15

10

Ser Leu Arg Leu Ser Cys Val Ala Ser Gly Asn Thr Phe Ser Ile Ile
20 25 30

15

Ala Met Ala Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Val Val
35 40 45

Ala Ser Ile Asn Ser Ile Gly Ser Thr Asn Tyr Ala Asp Ser Val Lys
50 55 60

20

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr Leu
65 70 75 80

25

Gln Met Ser Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

Ala Gly Asn Leu Leu Val Lys Arg Pro Tyr Trp Gly Gln Gly Thr Leu
100 105 110

30

Val Thr Val Ser Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Ala Ala
115 120 125

35

Ala His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser
130 135 140

Glu Glu Asp Leu Asn Gly Ala Ala
145 150

40

<210> 8

<211> 462

<212> DNA

45

<213> Artificial

<220>

50

<223> VHH with linker

<220>

<221> CDS

55

<222> (1)..(450)

EP 1 118 669 A2

5 <400> 8
 acc atg gcc cag gtg aaa ctg cag cag tct ggg gga gga ttg gtg cag 48
 Thr Met Ala Gln Val Lys Leu Gln Gln Ser Gly Gly Gly Leu Val Gln
 1 5 10 15

 gct ggg ggc cct ctg agg ctc tcc tgt gca gcc tct gga cgc acc ttc 96
 Ala Gly Gly Pro Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Phe
 20 25 30

 10 agt aac tat gcc gtg ggc tgg ttc cgc cag gct cca ggg aag gag cgt 144
 Ser Asn Tyr Ala Val Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg
 35 40 45

 gag ttt gtc gct gct att agc cgt gat ggt ggg cgc aca tac tat gcg 192
 Glu Phe Val Ala Ala Ile Ser Arg Asp Gly Gly Arg Thr Tyr Tyr Ala
 50 55 60
 15

 gac tcc gtg aag ggc cga ttc gcc gtc tcc aga gac tac gcc gag aac 240
 Asp Ser Val Lys Gly Arg Phe Ala Val Ser Arg Asp Tyr Ala Glu Asn
 65 70 75 80

 20 acg gtg tat ctg caa atg aac agc ctg aaa cct gag gac acg gcc gtt 288
 Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val
 85 90 95

 tat tac tgt aac aca agg gcc tac tgg ggc cag ggg acc cag gtc acc 336
 Tyr Tyr Cys Asn Thr Arg Ala Tyr Trp Gly Gln Gly Thr Gln Val Thr
 100 105 110
 25

 gtc tcc tca gcg cac cac agc gaa gac ccc agc tcc gcg gcc gcc cat 384
 Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala Ala Ala His
 115 120 125

 30 cac cat cac cat cac ggg gcc gca gaa caa aaa ctc atc tca gaa gag 432
 His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu
 130 135 140

 gat ctg aat ggg gcc gca tagtaacaat tg 462
 Asp Leu Asn Gly Ala Ala
 145 150
 35

 <210> 9

 <211> 150

 40 <212> PRT

 <213> Artificial

 45 <400> 9

 Thr Met Ala Gln Val Lys Leu Gln Gln Ser Gly Gly Gly Leu Val Gln
 1 5 10 15

 50 Ala Gly Gly Pro Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Phe
 20 25 30

 Ser Asn Tyr Ala Val Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg
 35 40 45
 55

EP 1 118 669 A2

	Glu Phe Val Ala Ala Ile Ser Arg Asp Gly Gly Arg Thr Tyr Tyr Ala	
	50 55 60	
5	Asp Ser Val Lys Gly Arg Phe Ala Val Ser Arg Asp Tyr Ala Glu Asn	
	65 70 75 80	
	Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val	
	85 90 95	
10	Tyr Tyr Cys Asn Thr Arg Ala Tyr Trp Gly Gln Gly Thr Gln Val Thr	
	100 105 110	
15	Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala Ala Ala His	
	115 120 125	
	His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu	
	130 135 140	
20	Asp Leu Asn Gly Ala Ala	
	145 150	
25	<210> 10	
	<211> 471	
	<212> DNA	
	<213> Artificial	
30		
	<220>	
	<223> VHH with linker	
35	<220>	
	<221> CDS	
	<222> (1)..(459)	
40		
	<400> 10	
	acc atg gcc cag gtg aaa ctg cag cag tct ggg gga gga ttg gtg cag	48
	Thr Met Ala Gln Val Lys Leu Gln Gln Ser Gly Gly Gly Leu Val Gln	
45	1 5 10 15	
	gct ggg ggc cct ctg agg ctc tcc tgt gca gcc tct gga cgc acc ttc	96
	Ala Gly Gly Pro Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Phe	
	20 25 30	
50	agt aac tat gcc gtg ggc tgg ttc cgc cag gct cca ggg aag gag cgt	144
	Ser Asn Tyr Ala Val Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg	
	35 40 45	
	gag ttt gtc gct gct att agc cgt gat ggt ggg cgc aca tac tat gcg	192
	Glu Phe Val Ala Ala Ile Ser Arg Asp Gly Gly Arg Thr Tyr Tyr Ala	
	50 55 60	
55	gac tcc gtg aag ggc cga ttc gcc gtc tcc aga gac tac gcc gag aac	240

EP 1 118 669 A2

	Asp Ser Val Lys Gly Arg Phe Ala Val Ser Arg Asp Tyr Ala Glu Asn	
	65 70 75 80	
5	acg gtg tat ctg caa atg aac agc ctg aaa cct gag gac acg gcc gtt Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val	288
	85 90 95	
	tat tac tgt aac aca agg gcc tac tgg ggc cag ggg acc cag gtc acc Tyr Tyr Cys Asn Thr Arg Ala Tyr Trp Gly Gln Gly Thr Gln Val Thr	336
	100 105 110	
10	gtc tcc tca gcg cac cac agc gaa gac ccc agc tcc gcg gcc gcc cat Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala Ala Ala His	384
	115 120 125	
	cac cat cac cat cac ggg gcc gca gaa caa aaa ctc atc tca gaa gag His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu	432
15	130 135 140	
	gat ctg aat agt gag aaa gat gag cta tgataacaat tg Asp Leu Asn Ser Glu Lys Asp Glu Leu	471
	145 150	
20	<210> 11	
	<211> 153	
	<212> PRT	
25	<213> Artificial	
	<400> 11	
30	Thr Met Ala Gln Val Lys Leu Gln Gln Ser Gly Gly Gly Leu Val Gln	
	1 5 10 15	
	Ala Gly Gly Pro Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Phe	
35	20 25 30	
	Ser Asn Tyr Ala Val Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg	
	35 40 45	
40	Glu Phe Val Ala Ala Ile Ser Arg Asp Gly Gly Arg Thr Tyr Tyr Ala	
	50 55 60	
	Asp Ser Val Lys Gly Arg Phe Ala Val Ser Arg Asp Tyr Ala Glu Asn	
45	65 70 75 80	
	Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val	
	85 90 95	
50	Tyr Tyr Cys Asn Thr Arg Ala Tyr Trp Gly Gln Gly Thr Gln Val Thr	
	100 105 110	
	Val Ser Ser Ala His His Ser Glu Asp Pro Ser Ser Ala Ala Ala His	
55	115 120 125	

EP 1 118 669 A2

His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu
 130 135 140

5 Asp Leu Asn Ser Glu Lys Asp Glu Leu
 145 150

<210> 12

10 <211> 38

<212> DNA

<213> Artificial

15 <220>

<223> synthetic insert

20 <400> 12
 agctgcgacg gcaagcttgg taccgggaat tctctaga 38

<210> 13

25 <211> 42

<212> DNA

<213> Artificial

30 <220>

<223> synthetic insert

35 <400> 13
 aatttctaga gaattcccg gaccaagctt gcttgcgacg gc 42

<210> 14

40 <211> 31

<212> DNA

<213> Artificial

45 <220>

<223> synthetic insert

50 <400> 14
 tcgaccccatg gcccgctagc caattggagc t 31

<210> 15

55 <211> 23

<212> DNA

EP 1 118 669 A2

<213> Artificial

5 <220>

<223> synthetic insert

<400> 15
ccaattggct agcgggccat ggg 23

10 <210> 16

<211> 22

15 <212> DNA

<213> Artificial

20 <220>

<223> primer

<400> 16
ccacccacga gggaacatcg tg 22

25 <210> 17

<211> 39

<212> DNA

30 <213> Artificial

35 <220>

<223> primer

<400> 17
gaattcccat ggtttacact cgaggtcctc tccaaatga 39

40 <210> 18

<211> 189

<212> DNA

45 <213> Artificial

50 <220>

<223> PCR product

<400> 18
ccacccacga gggaacatcg tggaaaaaga agacgttcca accacgtctt caaagcaagt 60
ggattgatgt gatatctcca ctgacgtaag ggatgacgca caatcccact atccttcgca 120
55 agacccttcc tttatataag gaagttcatt tcatttgag aggacctcga gtgtaaacca 180

	tggggaattc	189
5	<210> 19	
	<211> 21	
	<212> DNA	
10	<213> Artificial	
	<220>	
15	<223> sequencing primer	
	<400> 19	
	ccggcaacag gattcaatct t	21
20	<210> 20	
	<211> 40	
	<212> DNA	
25	<213> Artificial	
	<220>	
30	<223> synthetic insert	
	<400> 20	
	agctccatgg gatttgttct cttttcaciaa ttgccttcac	40
35	<210> 21	
	<211> 35	
	<212> DNA	
40	<213> Artificial	
	<220>	
45	<223> synthetic insert	
	<400> 21	
	ttcttcttgt ctctacactt ctcttattcc tagta	35
50	<210> 22	
	<211> 35	
	<212> DNA	
55	<213> Artificial	

EP 1 118 669 A2

<220>
 <223> synthetic insert
 5 <400> 22
 atatcccact cttgccgtgc ccaggtgcag ctgca 35
 <210> 23
 10 <211> 48
 <212> DNA
 <213> Artificial
 15 <220>
 <223> synthetic insert
 20 <400> 23
 gctgcacctg ggcacggcaa gagtgggata ttactaggaa taagagaa 48
 <210> 24
 <211> 54
 25 <212> DNA
 <213> Artificial
 30 <220>
 <223> synthetic insert
 35 <400> 24
 gtgtagagac aagaagaaat gaaggcaatt gtgaaaagag aacaaatccc atgg 54
 <210> 25
 <211> 24
 40 <212> DNA
 <213> Artificial
 45 <220>
 <223> synthetic insert
 50 <400> 25
 ggaaacagct atgaccatga ttac 24
 <210> 26
 <211> 20
 55 <212> DNA

EP 1 118 669 A2

<213> Artificial

 5 <220>
 <223> synthetic insert

 <400> 26
 tttcccagtc acgacgttgt 20

 10 <210> 27
 <211> 107

 15 <212> DNA
 <213> Artificial

 20 <220>
 <223> synthetic fragment encoding PRIa leader
 <220>
 <221> CDS
 25 <222> (3)..(104)

 30 <400> 27
 cc atg gga ttt gtt ctc ttt tca caa ttg cct tca ttt ctt ctt gtc 47
 Met Gly Phe Val Leu Phe Ser Gln Leu Pro Ser Phe Leu Leu Val
 1 5 10 15

 tct aca ctt ctc tta ttc cta gta ata tcc cac tct tgc cgt gcc cag 95
 Ser Thr Leu Leu Leu Phe Leu Val Ile Ser His Ser Cys Arg Ala Gln
 20 25 30
 35
 gtg cag ctg cag 107
 Val Gln Leu

 40 <210> 28
 <211> 34
 <212> PRT
 45 <213> Artificial

 50 <400> 28
 Met Gly Phe Val Leu Phe Ser Gln Leu Pro Ser Phe Leu Leu Val Ser
 1 5 10 15

 Thr Leu Leu Leu Phe Leu Val Ile Ser His Ser Cys Arg Ala Gln Val
 20 25 30
 55

5	<210> 29	
	<211> 21	
	<212> DNA	
10	<213> Artificial	
	<220>	
15	<223> PCR primer	
	<400> 29	
	gacatcccat ggcaagcatc a	21
20	<210> 30	
	<211> 21	
	<212> DNA	
25	<213> Artificial	
	<220>	
30	<223> PCR primer	
	<400> 30	
	aagcttggtta acagccctta a	21
35	<210> 31	
	<211> 21	
	<212> DNA	
40	<213> Artificial	
	<220>	
45	<223> PCR primer	
	<400> 31	
	agggttggtta acaaacttga t	21
50	<210> 32	
	<211> 44	
	<212> DNA	
55	<213> Artificial	

EP 1 118 669 A2

<220>

<223> PCR primer

5 <400> 32
agactgctgc agctgcacct gctttccaca aacaatggta gctg 44

<210> 33

10 <211> 19
<212> DNA
<213> Artificial

15 <220>

<223> sequencing primer

20 <400> 33
aattaaccct cactaaagg 19

<210> 34

25 <211> 254
<212> DNA
<213> Artificial

30 <220>

<223> GBSS leader

35 <220>

<221> CDS

<222> (3)..(254)

40 <400> 34
cc atg gca agc atc aca gct tca cac cac ttt gtg tca aga agc caa 47
Met Ala Ser Ile Thr Ala Ser His His Phe Val Ser Arg Ser Gln
1 5 10 15

45 act tca cta gac acc aaa tca acc ttg tca cag ata gga ctc agg aac 95
Thr Ser Leu Asp Thr Lys Ser Thr Leu Ser Gln Ile Gly Leu Arg Asn
20 25 30

50 cat act ctg act cac aat ggt tta agg gct gtt aac aaa ctt gat ggg 143
His Thr Leu Thr His Asn Gly Leu Arg Ala Val Asn Lys Leu Asp Gly
35 40 45

55 ctc caa tca aga act aat act aag gta aca ccc aag atg gca tcc aga 191
Leu Gln Ser Arg Thr Asn Thr Lys Val Thr Pro Lys Met Ala Ser Arg
50 55 60

65 act gag acc aag aga cct gga tgc tca gct acc att gtt tgt gga aaa 239
Thr Glu Thr Lys Arg Pro Gly Cys Ser Ala Thr Ile Val Cys Gly Lys
65 70 75

cag gtg cag ctg cag
Gln Val Gln Leu Gln
80

254

5

<210> 35

<211> 84

10

<212> PRT

<213> Artificial

15

<400> 35

Met Ala Ser Ile Thr Ala Ser His His Phe Val Ser Arg Ser Gln Thr
1 5 10 15

20

Ser Leu Asp Thr Lys Ser Thr Leu Ser Gln Ile Gly Leu Arg Asn His
20 25 30

Thr Leu Thr His Asn Gly Leu Arg Ala Val Asn Lys Leu Asp Gly Leu
35 40 45

25

Gln Ser Arg Thr Asn Thr Lys Val Thr Pro Lys Met Ala Ser Arg Thr
50 55 60

30

Glu Thr Lys Arg Pro Gly Cys Ser Ala Thr Ile Val Cys Gly Lys Gln
65 70 75 80

Val Gln Leu Gln

35

<210> 36

<211> 18

<212> DNA

40

<213> Artificial

<220>

45

<223> synthetic insert

<400> 36
catgcaggtg cagctgca

18

50

<210> 37

<211> 10

<212> DNA

55

<213> Artificial

<220>
 5 <223> synthetic insert
 <400> 37
 gctgcacctg 10
 <210> 38
 10 <211> 24
 <212> DNA
 <213> Artificial
 15
 <220>
 <223> PCR primer
 20 <400> 38
 cgcaagaccc ttcctctata taag 24
 <210> 39
 25 <211> 60
 <212> DNA
 <213> Artificial
 30
 <220>
 <223> PCR primer
 35 <400> 39
 gagctcgaat tottattata gctcatcttt ctctgaattc agatcctctt ctgagatgag 60
 <210> 40
 40 <211> 26
 <212> PRT
 <213> Artificial
 45
 <220>
 <223> linker
 50 <400> 40
 Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15
 Gly Gly Gly Gly Ser Asp Ile Glu Leu Thr
 20 25
 55

EP 1 118 669 A2

	<210> 41	
	<211> 20	
5	<212> DNA	
	<213> Artificial	
10	<220>	
	<223> PCR primer	
	<400> 41	
	atcctcaact tccaatcaga	20
15	<210> 42	
	<211> 19	
	<212> DNA	
20	<213> Artificial	
25	<220>	
	<223> PCR primer	
	<400> 42	
	ttcttgagag atagcttga	19
30	<210> 43	
	<211> 30	
	<212> DNA	
35	<213> Artificial	
40	<220>	
	<223> synthetic insert	
	<400> 43	
	gatcccatgg cccgctagcc aattggagct	30
45	<210> 44	
	<211> 22	
	<212> DNA	
50	<213> Artificial	
55	<220>	
	<223> synthetic insert	

EP 1 118 669 A2

	<400> 44	
	ccaattggct agcgggccat gg	22
5	<210> 45	
	<211> 29	
	<212> DNA	
10	<213> Artificial	
	<220>	
15	<223> synthetic insert	
	<400> 45	
	gatccacctc gagtgtaaac catggcccg	29
20	<210> 46	
	<211> 29	
	<212> DNA	
25	<213> Artificial	
	<220>	
30	<223> synthetic insert	
	<400> 46	
	ctagcggggc atggtttaca ctcgaggtg	29
35	<210> 47	
	<211> 20	
	<212> DNA	
40	<213> Artificial	
	<220>	
45	<223> PCR primer	
	<400> 47	
	attgcctacg gcagccgctg	20
50	<210> 48	
	<211> 51	
	<212> DNA	
55	<213> Artificial	

EP 1 118 669 A2

<220>
 <223> PCR primer
 5 <400> 48
 tccaaccaat tggtactatg cggccccatt cagatcctct tctgagatga g 51
 <210> 49
 10 <211> 25
 <212> DNA
 <213> Artificial
 15 <220>
 <223> sequencing primer
 20 <400> 49
 gtctgtctaa agtaaagtag atgcg 25
 <210> 50
 <211> 60
 25 <212> DNA
 <213> Artificial
 30 <220>
 <223> PCR primer
 <400> 50
 35 tccaaccaat tggtatcata gctcatcttt ctactattc agatcctctt ctgagatgag 60
 <210> 51
 <211> 29
 40 <212> DNA
 <213> Artificial
 45 <220>
 <223> PCR primer
 <400> 51
 50 agtcccccat ggtacgtcct gtagaaacc 29
 <210> 52
 <211> 25
 55 <212> DNA

EP 1 118 669 A2

<213> Artificial

 5 <220>
 <223> PCR primer
 <400> 52
 cgttttcgtc ggtaatcacc attcc 25
 10
 <210> 53
 <211> 24
 <212> DNA
 15 <213> Artificial

 <220>
 20 <223> sequencing primer
 <400> 53
 cgcaagaccc ttcctttata taag 24
 25
 <210> 54
 <211> 1154
 <212> DNA
 30 <213> Artificial

 <220>
 35 <223> HCV33-hinge-CH2-CH3
 <220>
 <221> CDS
 40 <222> (3)..(1136)

 <400> 54
 cc atg gag gtg cag ctg cag gag tca ggg gga gga ttg gtg cag gct 47
 45 Met Glu Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala
 1 5 10 15
 ggg ggc tct ctg aga ctc tcc tgt gca gcc tgg gga cgc gcc acc agt 95
 Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser
 20 25 30
 50 ggt cat ggt cac tat ggt atg ggc tgg ttc cgc cag gtt cca ggg aag 143
 Gly His Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys
 35 40 45
 gag cgt gag ttt gtc gca gct att agg tgg agt ggt aaa gag aca tgg 191
 55 Glu Arg Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp
 50 55 60

EP 1 118 669 A2

	tat aaa gac tcc gtg aag ggc cga ttc acc atc tcc aga gat aac gcc	239
	Tyr Lys Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala	
	65 70 75	
5	aag act acg gtt tat ctg caa atg aac agc ctg aaa cct gaa gat acg	287
	Lys Thr Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr	
	80 85 90 95	
	gcc gtt tat tat tgt gcc gct cga ccg gtc cgc gtg gat gat att tcc	335
10	Ala Val Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser	
	100 105 110	
	ctg ccg gtt ggg ttt gac tac tgg ggc cag ggg acc cag gtc acc gtc	383
	Leu Pro Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val	
	115 120 125	
15	tcc tca gaa ccc aag aca cca aaa cca caa cca caa cca caa cca caa	431
	Ser Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Gln Pro Gln Pro Gln	
	130 135 140	
	cca caa ccc aat cct aca aca gaa tcc aag tgt ccc aaa tgt cca gcc	479
20	Pro Gln Pro Asn Pro Thr Thr Glu Ser Lys Cys Pro Lys Cys Pro Ala	
	145 150 155	
	cct gag ctc ctg gga ggg ccc tca gtc ttc atc ttc ccc ccg aaa ccc	527
	Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Ile Phe Pro Pro Lys Pro	
	160 165 170 175	
25	aag gac gtc ctc tcc att tct ggg agg ccc gag gtc acg tgc gtt gtg	575
	Lys Asp Val Leu Ser Ile Ser Gly Arg Pro Glu Val Thr Cys Val Val	
	180 185 190	
	gta gac gtg ggc cag gaa gac ccc gag gtc agt ttc aac tgg tac att	623
30	Val Asp Val Gly Gln Glu Asp Pro Glu Val Ser Phe Asn Trp Tyr Ile	
	195 200 205	
	gat ggc gca gag gtg cga acg gcc aac acg agg cca aaa gag gaa cag	671
	Asp Gly Ala Glu Val Arg Thr Ala Asn Thr Arg Pro Lys Glu Glu Gln	
	210 215 220	
35	ttc aac agc acg tac cgc gtg gtc agc gtc ctg ccc atc cag cac cag	719
	Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Pro Ile Gln His Gln	
	225 230 235	
	gac tgg ctg acg ggg aaa gag ttc aaa tgc aag gtc aac aac aaa gct	767
40	Asp Trp Leu Thr Gly Lys Glu Phe Lys Cys Lys Val Asn Asn Lys Ala	
	240 245 250 255	
	ctc ccg gcc ccc atc gag aag acc atc tcc aag gcc aaa ggg cag acc	815
	Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Thr	
	260 265 270	
45	cgg gag ccg cag gtg tac gcc ctg gcc cca cac cgg gaa gag ctg gcc	863
	Arg Glu Pro Gln Val Tyr Ala Leu Ala Pro His Arg Glu Glu Leu Ala	
	275 280 285	
	aag gac acc gtg agc gta acc tgc ctg gtc aaa ggc ttc tac cca cct	911
50	Lys Asp Thr Val Ser Val Thr Cys Leu Val Lys Gly Phe Tyr Pro Pro	
	290 295 300	
	gat atc aac gtt gag tgg cag agg aac ggt cag ccg gag tca gag gcc	959
	Asp Ile Asn Val Glu Trp Gln Arg Asn Gly Gln Pro Glu Ser Glu Gly	
	305 310 315	
55	acc tac gcc acc acg cca ccc cag ctg gac aac gac ggg acc tac ttc	1007
	Thr Tyr Ala Thr Thr Pro Pro Gln Leu Asp Asn Asp Gly Thr Tyr Phe	
	320 325 330 335	

EP 1 118 669 A2

	ctc tac agc aag ctc tcg gtg gga aag aac acg tgg cag cgg gga gaa	1055
	Leu Tyr Ser Lys Leu Ser Val Gly Lys Asn Thr Trp Gln Arg Gly Glu	
	340 345 350	
5	acc ttc acc tgt gtg gtg atg cac gag gcc ctg cac aac cac tac acc	1103
	Thr Phe Thr Cys Val Val Met His Glu Ala Leu His Asn His Tyr Thr	
	355 360 365	
10	cag aaa tcc atc acc cag tct tcg ggt aaa taa taagaattcg agctcgaa	1154
	Gln Lys Ser Ile Thr Gln Ser Ser Gly Lys	
	370 375	
	<210> 55	
15	<211> 377	
	<212> PRT	
	<213> Artificial	
20	<400> 55	
	Met Glu Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly	
	1 5 10 15	
25	Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly	
	20 25 30	
30	His Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu	
	35 40 45	
	Arg Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr	
	50 55 60	
35	Lys Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys	
	65 70 75 80	
40	Thr Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala	
	85 90 95	
	Val Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu	
	100 105 110	
45	Pro Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser	
	115 120 125	
50	Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Gln Pro Gln Pro Gln Pro	
	130 135 140	
	Gln Pro Asn Pro Thr Thr Glu Ser Lys Cys Pro Lys Cys Pro Ala Pro	
	145 150 155 160	
55	Glu Leu Leu Gly Gly Pro Ser Val Phe Ile Phe Pro Pro Lys Pro Lys	
	165 170 175	

EP 1 118 669 A2

Asp Val Leu Ser Ile Ser Gly Arg Pro Glu Val Thr Cys Val Val Val
 180 185 190
 5
 Asp Val Gly Gln Glu Asp Pro Glu Val Ser Phe Asn Trp Tyr Ile Asp
 195 200 205
 10
 Gly Ala Glu Val Arg Thr Ala Asn Thr Arg Pro Lys Glu Glu Gln Phe
 210 215 220
 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Pro Ile Gln His Gln Asp
 225 230 235 240
 15
 Trp Leu Thr Gly Lys Glu Phe Lys Cys Lys Val Asn Asn Lys Ala Leu
 245 250 255
 20
 Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Thr Arg
 260 265 270
 Glu Pro Gln Val Tyr Ala Leu Ala Pro His Arg Glu Glu Leu Ala Lys
 275 280 285
 25
 Asp Thr Val Ser Val Thr Cys Leu Val Lys Gly Phe Tyr Pro Pro Asp
 290 295 300
 30
 Ile Asn Val Glu Trp Gln Arg Asn Gly Gln Pro Glu Ser Glu Gly Thr
 305 310 315 320
 Tyr Ala Thr Thr Pro Pro Gln Leu Asp Asn Asp Gly Thr Tyr Phe Leu
 325 330 335
 35
 Tyr Ser Lys Leu Ser Val Gly Lys Asn Thr Trp Gln Arg Gly Glu Thr
 340 345 350
 40
 Phe Thr Cys Val Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
 355 360 365
 Lys Ser Ile Thr Gln Ser Ser Gly Lys
 370 375
 45
 <210> 56
 <211> 1172
 <212> DNA
 50
 <213> Artificial
 <220>
 55
 <223> hinge-HCV33-CH2-CH3-SEKDEL

EP 1 118 669 A2

<220>

<221> CDS

5 <222> (3)..(1154)

<400> 56

10	cc atg gag gtg cag ctg cag gag tca ggg gga gga ttg gtg cag gct	47
	Met Glu Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala	
	1 5 10 15	
	ggg ggc tct ctg aga ctc tcc tgt gca gcc tcg gga cgc gcc acc agt	95
	Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser	
	20 25 30	
15	ggg cat ggt cac tat ggt atg ggc tgg ttc cgc cag gtt cca ggg aag	143
	Gly His Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys	
	35 40 45	
	gag cgt gag ttt gtc gca gct att agg tgg agt ggt aaa gag aca tgg	191
	Glu Arg Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp	
20	50 55 60	
	tat aaa gac tcc gtg aag ggc cga ttc acc atc tcc aga gat aac gcc	239
	Tyr Lys Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala	
	65 70 75	
25	aag act acg gtt tat ctg caa atg aac agc ctg aaa cct gaa gat acg	287
	Lys Thr Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr	
	80 85 90 95	
	gcc gtt tat tat tgt gcc gct cga ccg gtc cgc gtg gat gat att tcc	335
	Ala Val Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser	
30	100 105 110	
	ctg ccg gtt ggg ttt gac tac tgg ggc cag ggg acc cag gtc acc gtc	383
	Leu Pro Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val	
	115 120 125	
35	tcc tca gaa ccc aag aca cca aaa cca caa cca caa cca caa cca	431
	Ser Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Gln Pro Gln Pro Gln	
	130 135 140	
	cca caa ccc aat cct aca aca gaa tcc aag tgt ccc aaa tgt cca gcc	479
	Pro Gln Pro Asn Pro Thr Thr Glu Ser Lys Cys Pro Lys Cys Pro Ala	
40	145 150 155	
	cct gag ctc ctg gga ggg ccc tca gtc ttc atc ttc ccc ccg aaa ccc	527
	Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Ile Phe Pro Pro Lys Pro	
	160 165 170 175	
45	aag gac gtc ctc tcc att tct ggg agg ccc gag gtc acg tgc gtt gtg	575
	Lys Asp Val Leu Ser Ile Ser Gly Arg Pro Glu Val Thr Cys Val Val	
	180 185 190	
	gta gac gtg ggc cag gaa gac ccc gag gtc agt ttc aac tgg tac att	623
	Val Asp Val Gly Gln Glu Asp Pro Glu Val Ser Phe Asn Trp Tyr Ile	
50	195 200 205	
	gat ggc gca gag gtg cga acg gcc aac acg agg cca aaa gag gaa cag	671
	Asp Gly Ala Glu Val Arg Thr Ala Asn Thr Arg Pro Lys Glu Glu Gln	
	210 215 220	
55	ttc aac agc acg tac cgc gtg gtc agc gtc ctg ccc atc cag cac cag	719
	Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Pro Ile Gln His Gln	
	225 230 235	

EP 1 118 669 A2

	gac tgg ctg acg ggg aaa gag ttc aaa tgc aag gtc aac aac aaa gct	767
	Asp Trp Leu Thr Gly Lys Glu Phe Lys Cys Lys Val Asn Asn Lys Ala	
	240 245 250 255	
5	ctc ccg gcc ccc atc gag aag acc atc tcc aag gcc aaa ggg cag acc	815
	Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Thr	
	260 265 270	
10	cgg gag ccg cag gtg tac gcc ctg gcc cca cac cgg gaa gag ctg gcc	863
	Arg Glu Pro Gln Val Tyr Ala Leu Ala Pro His Arg Glu Glu Leu Ala	
	275 280 285	
	aag gac acc gtg agc gta acc tgc ctg gtc aaa ggc ttc tac cca cct	911
	Lys Asp Thr Val Ser Val Thr Cys Leu Val Lys Gly Phe Tyr Pro Pro	
	290 295 300	
15	gat atc aac gtt gag tgg cag agg aac ggt cag ccg gag tca gag ggc	959
	Asp Ile Asn Val Glu Trp Gln Arg Asn Gly Gln Pro Glu Ser Glu Gly	
	305 310 315	
20	acc tac gcc acc acg cca ccc cag ctg gac aac gac ggg acc tac ttc	1007
	Thr Tyr Ala Thr Thr Pro Pro Gln Leu Asp Asn Asp Gly Thr Tyr Phe	
	320 325 330 335	
	ctc tac agc aag ctc tcg gtg gga aag aac acg tgg cag ccg gga gaa	1055
	Leu Tyr Ser Lys Leu Ser Val Gly Lys Asn Thr Trp Gln Arg Gly Glu	
	340 345 350	
25	acc ttc acc tgt gtg gtg atg cac gag gcc ctg cac aac cac tac acc	1103
	Thr Phe Thr Cys Val Val Met His Glu Ala Leu His Asn His Tyr Thr	
	355 360 365	
30	cag aaa tcc atc acc cag tct tcg ggt aaa tct gag aaa gat gag cta	1151
	Gln Lys Ser Ile Thr Gln Ser Ser Gly Lys Ser Glu Lys Asp Glu Leu	
	370 375 380	
	taa taagaattcg agctcgaa	1172
35	<210> 57	
	<211> 383	
	<212> PRT	
	<213> Artificial	
40	<400> 57	
45	Met Glu Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly	
	1 5 10 15	
	Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly	
	20 25 30	
50	His Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu	
	35 40 45	
55	Arg Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr	
	50 55 60	

EP 1 118 669 A2

	Lys	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	
	65					70					75					80	
5	Thr	Thr	Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Lys	Pro	Glu	Asp	Thr	Ala	
					85					90					95		
	Val	Tyr	Tyr	Cys	Ala	Ala	Arg	Pro	Val	Arg	Val	Asp	Asp	Ile	Ser	Leu	
				100					105					110			
10	Pro	Val	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	
			115					120					125				
	Ser	Glu	Pro	Lys	Thr	Pro	Lys	Pro	Gln	Pro	Gln	Pro	Gln	Pro	Gln	Pro	
15		130					135					140					
	Gln	Pro	Asn	Pro	Thr	Thr	Glu	Ser	Lys	Cys	Pro	Lys	Cys	Pro	Ala	Pro	
	145					150					155					160	
20	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Lys	Pro	Lys	
				165						170					175		
	Asp	Val	Leu	Ser	Ile	Ser	Gly	Arg	Pro	Glu	Val	Thr	Cys	Val	Val	Val	
25				180					185					190			
	Asp	Val	Gly	Gln	Glu	Asp	Pro	Glu	Val	Ser	Phe	Asn	Trp	Tyr	Ile	Asp	
			195					200					205				
30	Gly	Ala	Glu	Val	Arg	Thr	Ala	Asn	Thr	Arg	Pro	Lys	Glu	Glu	Gln	Phe	
		210					215					220					
	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Pro	Ile	Gln	His	Gln	Asp	
35		225				230					235					240	
	Trp	Leu	Thr	Gly	Lys	Glu	Phe	Lys	Cys	Lys	Val	Asn	Asn	Lys	Ala	Leu	
				245						250					255		
40	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Thr	Arg	
				260					265					270			
	Glu	Pro	Gln	Val	Tyr	Ala	Leu	Ala	Pro	His	Arg	Glu	Glu	Leu	Ala	Lys	
45			275					280					285				
	Asp	Thr	Val	Ser	Val	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Pro	Asp	
		290					295					300					
50	Ile	Asn	Val	Glu	Trp	Gln	Arg	Asn	Gly	Gln	Pro	Glu	Ser	Glu	Gly	Thr	
	305					310					315					320	
	Tyr	Ala	Thr	Thr	Pro	Pro	Gln	Leu	Asp	Asn	Asp	Gly	Thr	Tyr	Phe	Leu	
55					325					330					335		

EP 1 118 669 A2

	Tyr Ser Lys Leu Ser Val Gly Lys Asn Thr Trp Gln Arg Gly Glu Thr	
	340 345 350	
5	Phe Thr Cys Val Val Met His Glu Ala Leu His Asn His Tyr Thr Gln	
	355 360 365	
	Lys Ser Ile Thr Gln Ser Ser Gly Lys Ser Glu Lys Asp Glu Leu	
	370 375 380	
10	<210> 58	
	<211> 31	
15	<212> DNA	
	<213> Artificial	
20	<220>	
	<223> PCR primer	
	<400> 58	
	aggggaccca ggtcacgctc tcttcagaac c	31
25	<210> 59	
	<211> 37	
	<212> DNA	
30	<213> Artificial	
35	<220>	
	<223> PCR primer	
	<400> 59	
	gagctttgtt gttgaccttg catttgaact ctttccc	37
40	<210> 60	
	<211> 26	
	<212> DNA	
45	<213> Artificial	
50	<220>	
	<223> PCR primer	
	<400> 60	
	caaatgcaag gtcaacaaca aagctc	26
55	<210> 61	

EP 1 118 669 A2

<211> 42
 <212> DNA
 5 <213> Artificial

 <220>
 10 <223> PCR primer
 <400> 61
 ttcgagctcg aattcttatt atttaccga agactgggtg at 42

 15 <210> 62
 <211> 27
 <212> DNA
 20 <213> Artificial

 <220>
 25 <223> sequencing primer
 <400> 62
 ctgaggagac ggtgacctgg gtccct 27

 30 <210> 63
 <211> 46
 <212> DNA
 35 <213> Artificial

 <220>
 40 <223> PCR primer
 <400> 63
 agccctgag ctctgggag ggccctcagt ctcatcttc ccccg 46

 45 <210> 64
 <211> 61
 <212> DNA
 50 <213> Artificial

 <220>
 55 <223> PCR primer
 <400> 64
 ttcgagctcg aattcttatt atttaccga agactgggtg atggatttct gggtagtg 60

g 61

5 <210> 65
 <211> 79
 <212> DNA
 10 <213> Artificial

<220>
 15 <223> PCR primer
 <400> 65
 ttcgagctcg aattcttatt atagctcatc tttctcagat ttacccgaag actgggtgat 60
 ggatttctgg gtgtagtgg 79

20 <210> 66
 <211> 461
 <212> DNA
 25 <213> Artificial

<220>
 30 <223> VHH with linker
 <220>
 <221> CDS
 35 <222> (3)..(449)

<400> 66
 40 cc atg gcc cag gtg cag ctg cag gag tct ggg gga ggc ttg gtg cag 47
 Met Ala Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln
 1 5 10 15
 gct ggg ggg tct ctg agg ctc tcc tgt gca gcc tct gga agc att ttc 95
 Ala Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Ser Ile Phe
 20 25 30
 45 aga cgt ccg cat atg ggt tgg ttc cgc cag gct cca ggg cag gag cgc 143
 Arg Arg Pro His Met Gly Trp Phe Arg Gln Ala Pro Gly Gln Glu Arg
 35 40 45
 gag ttg gtc gca ctg att tct gcg ggt ggt cgt aca tgg tat gca gac 191
 Glu Leu Val Ala Leu Ile Ser Ala Gly Gly Arg Thr Trp Tyr Ala Asp
 50 55 60
 tcc gtg aag ggc cga ttc acc atc tcc aga gac aac gcc aag aac acg 239
 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr
 65 70 75
 55 ctg tat ctg caa atg aac agc ctg aaa cct gag gac acg gcc gtt tat 287

EP 1 118 669 A2

	Leu Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr	
	80 85 90 95	
5	tac tgt act gcc ggg ggt tgc tac tgg ggc cag ggg acc cag gtc acc Tyr Cys Thr Ala Gly Gly Ser Tyr Trp Gly Gln Gly Thr Gln Val Thr	335
	100 105 110	
	gtc gcc tca gaa ccc aag aca cca aaa cca caa cca gcg gcc gcc cat Val Ala Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Ala Ala Ala His	383
	115 120 125	
10	cac cat cac cat cac ggg gcc gca gaa caa aaa ctc atc tca gaa gag His His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu	431
	130 135 140	
15	gat ctg aat ggg gcc gca tagtaacaat tg Asp Leu Asn Gly Ala Ala	461
	145	
	<210> 67	
20	<211> 149	
	<212> PRT	
	<213> Artificial	
25	<400> 67	
	Met Ala Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala	
	1 5 10 15	
30	Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Ser Ile Phe Arg	
	20 25 30	
35	Arg Pro His Met Gly Trp Phe Arg Gln Ala Pro Gly Gln Glu Arg Glu	
	35 40 45	
	Leu Val Ala Leu Ile Ser Ala Gly Gly Arg Thr Trp Tyr Ala Asp Ser	
	50 55 60	
40	Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu	
	65 70 75 80	
45	Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr	
	85 90 95	
	Cys Thr Ala Gly Gly Ser Tyr Trp Gly Gln Gly Thr Gln Val Thr Val	
	100 105 110	
50	Ala Ser Glu Pro Lys Thr Pro Lys Pro Gln Pro Ala Ala Ala His His	
	115 120 125	
55	His His His His Gly Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu Asp	
	130 135 140	

Leu Asn Gly Ala Ala
145

5

Claims

- 10 1. A method for modifying a plant to produce an antibody or an active fragment or derivative thereof, or a protein functionally equivalent thereto, in a desired cellular compartment comprising introducing into a plant a DNA sequence encoding a heavy chain immunoglobulin or an active fragment or derivative thereof, or a sequence encoding a protein functionally equivalent thereto, said DNA sequence being operably linked to one or more promoters and provided, as appropriate, with an additional sequence encoding a peptide sequence capable of targeting said antibody or fragment or derivative thereof to said desired cellular compartment.
- 15 2. A method according to claim 1 wherein the DNA sequence encoding the heavy chain immunoglobulin or fragment or derivative thereof is obtainable from camelids.
- 20 3. A method according to claim 1 or claim 2 wherein the plant is selected from tobacco, pea, potato, spinach, tomato or tea.
4. A method according to any one of claims 1 to 3 wherein the heavy chain immunoglobulin or active fragment or derivative thereof binds to a protein present in the plant.
- 25 5. A method according to any one of claims 1 to 3 wherein the heavy chain immunoglobulin or active fragment or derivative thereof binds to a plant or animal pathogen.
- 30 6. A method according to any one of claims 1 to 4 wherein the heavy chain immunoglobulin or active fragment or derivative thereof binds to a plant hormone or metabolite.
7. A plant prepared according to the method of any one of claims 1 to 6.
- 35 8. A modified plant having, in a desired cellular compartment, enhanced levels of heavy chain immunoglobulins or active fragments or derivatives thereof or proteins functionally equivalent thereto compared to equivalent but unmodified plants.
9. Seeds, fruits, progeny and hybrids of a plant according to claim 7 or 8.
- 40 10. A food product comprising a plant according to claim 7 or 8.
11. Use of a DNA sequence encoding a heavy chain immunoglobulin or an active fragment or derivative thereof or one or more sequences encoding a protein functionally equivalent thereto to cause a plant to produce said immunoglobulin or fragment or derivative thereof, or protein functionally equivalent thereto, in a desired cellular compartment.
- 45 12. Use according to claim 11 to increase pathogen resistance in a plant.
13. Use according to claim 11 to modulate plant metabolism.
- 50 14. A method for preparing a heavy chain immunoglobulin or an active fragment or derivative thereof comprising the steps of:
 - (i) modifying a plant according to the method of any of claims 1 to 6, and
 - 55 (ii) extracting from said modified plant the heavy chain immunoglobulin or active fragment or derivative thereof produced therein.

Fig.1.

*Pst*I

CAGGTGCAGCTGCAGGAGTCAGGGGGAGGATTGGTGCAGGCTGGGGGCTCTCTGAGACTC
 Q V Q L Q E S G G G L V Q A G G S L R L

TCCTGTGCAGCCTCGGGACGCGCCACCAGTGGTCATGGTCACTATGGTATGGGCTGGTTTC
 S C A A S G R A T S G H G H Y G M G W F

CGCCAGGTTCCAGGGAAGGAGCGTGAGTTTGTCGCAGCTATTAGGTGGAGTGGTAAAGAG
 R Q V P G K E R E F V A A I R W S G K E

ACATGGTATAAAGACTCCGTGAAGGGCCGATTCACCATCTCCAGAGATAACGCCAAGACT
 T W Y K D S V K G R F T I S R D N A K T

ACGGTTTATCTGCAAATGAACAGCCTGAAACCTGAAGATACGGCCGTTTATTATTGTGCC
 T V Y L Q M N S L K P E D T A V Y Y C A

GCTCGACCGGTCCGCGTGGATGATATTTCCCTGCCGTTGGGTTTGACTIONTGGGGCCAG
 A R P V R V D D I S L P V G F D Y W G Q

GGGACCCAGGTCACCGTCTCCTCAGAACAAAACTCATCTCAGAAGAGGATCTGAATTAA
 G T Q V T V S S E Q K L I S E E D L N

TAAGGGCTAAGCTCGAATTC
 EcoRI

Fig.2A.




Fig.2B.




Fig.3A.




Fig.3B.




Fig.4.




Fig.5.




Fig.6.

*Pst*I
 CAGGTGCAGCTGCAGCAGTCAGGGGAGGCTTGGTGCAGGCTGGGGGGTCTCTGAGACTC
 Q V Q L Q Q S G G G L V Q A G G S L R L
 TCCTGTGTAGCTTCTGAAAGCAGCTTCAGCAACAATCACATGGGCTGGTACCGCCGGGCT
 S C V A S E S S F S N N H M G W Y R R A
 CCAGGGAACCAGCGCGAGCTGGTCGCAACTATTAGTCCTGGTGGTAGCACACACTATGTA
 P G N Q R E L V A T I S P G G S T H Y V
 GACTCCGTGAAGGGCCGATTCACCATCTCCCGAGACAACGCCAAGAACACAGTGTATCTA
 D S V K G R F T I S R D N A K N T V Y L
 CAAATGGACAGCCTGAAACCAGAGGACACGGCCGTCTATTACTGTGCTGCCAAGGGGAGG
 Q M D S L K P E D T A V Y Y C A A K G R

*Pst*I
 GGGCTGCAGGCTATGCAGTACTGGGGCCAGGGGACCCTGGTCACCGTCTCCTCAGCGCAC
 G L Q A M Q Y W G Q G T L V T V S S A H
 CACAGCGAAGACCCAGCTCCGCGGGCCGCCATCACCATCACCATCACGGGGCCGCAGAA
 H S E D P S S A A A H H H H H G A A E
 CAAAACTCATCTCAGAAGAGGATCTGAATGGGGCCGCATAGTAACAATTG
 Q K L I S E E D L N G A A *Mun*I

Fig.7.




Fig.8.





Fig.9.




Fig.10A.




Fig.10B.




Fig.11.





Fig.12.

*Pst*I
 CAGGTGCAGCTGCAGGAGTCTGGGGGAGGCCTGGTGCAGGCTGGGGGGTCTCTGAGACTC
 Q V Q L Q E S G G G L V Q A G G S L R L
 TCCTGTGTAGCCTCTGGAAACACCTTCAGTATCATAGCTATGGCCTGGTACCGCCAGGCT
 S C V A S G N T F S I I A M A W Y R Q A
 CCAGGGAAGCAGCGCGAGGTGGTCGCAAGTATTAATAGTATTGGCAGCACAAATTATGCA
 P G K Q R E V V A S I N S I G S T N Y A
 GACTCCGTGAAGGGGCGATTACCATCTCCAGAGACAACGCCAAGAACACAGTGTATCTG
 D S V K G R F T I S R D N A K N T V Y L
 CAAATGAGCAGCCTGAAACCTGAGGACACGGCCGTCTATTACTGTGCTGCCGGTAATTTG
 Q M S S L K P E D T A V Y Y C A A G N L
 CTGGTTAAGAGGCCTTACTGGGGCCAGGGGACCCTGGTCACCGTCTCCTCAGAACCCAAG
 L V K R P Y W G Q G T L V T V S S E P K
 ACACCAAACCACAACCAGCGGCCGCCCATCACCATCACCATCACGGGGCCGCAGAACAA
 T P K P Q P A A A H H H H H H G A A E Q
 AAACATCTCAGAAGAGGATCTGAATGGGGCCGCATAGTAACAATTG
 K L I S E E D L N G A A *Mun*I

Fig.13.




Fig.14.

NcoI *PstI*
ACCATGGCCCAGGTGAAACCTGCAGCAGTCTGGGGGAGGATTGGTGCAGGCTGGGGGCCCT
 T M A Q V K L Q Q S G G G L V Q A G G P
 CTGAGGCTCTCCTCTGCAGCCTCTGGACGCACCTTCAGTAACTATGCCGTGGGCTGGTTC
 L R L S C A A S G R T F S N Y A V G W F
 CGCCAGGCTCCAGGGAAGGAGCGTGAGTTTGTGCTGCTATTAGCCGTGATGGTGGGCGC
 R Q A P G K E R E F V A A I S R D G G R
 ACATACTATGCGGACTCCGTGAAGGGCCGATTCGCCGTCTCCAGAGACTACGCCGAGAAC
 T Y Y A D S V K G R F A V S R D Y A E N
 ACGGTGTATCTGCAAATGAACAGCCTGAAACCTGAGGACACGGCCGTTTATTACTGTAAC
 T V Y L Q M N S L K P E D T A V Y Y C N
 ACAAGGGCCTACTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCAGCGCACCACAGCGAA
 T R A Y W G Q G T Q V T V S S A H H S E
 GACCCAGCTCCGCGGCCGCCCATCACCATCACCATCACGGGGCCGCAGAACAAAACTC
 D P S S A A A H H H H H H G A A E Q K L
 ATCTCAGAAGAGGATCTGAATGGGGCCGCATAGTAACCAATTG
 I S E E D L N G A A *MunI*

Fig.15.

NcoI *PstI*
ACCATGGCCCAGGTGAAACCTGCAGCAGTCTGGGGGAGGATTGGTGCAGGCTGGGGGCCCT
 T M A Q V K L Q Q S G G G L V Q A G G P
 CTGAGGCTCTCCTGTGCAGCCTCTGGACGCACCTTCAGTAACTATGCCGTGGGCTGGTTC
 L R L S C A A S G R T F S N Y A V G W F
 CGCCAGGCTCCAGGGAAGGAGCGTGAGTTTGTGCTGCTATTAGCCGTGATGGTGGGCGC
 R Q A P G K E R E F V A A I S R D G G R
 ACATACTATGCGGACTCCGTGAAGGGCCGATTCGCCGTCTCCAGAGACTACGCCGAGAAC
 T Y Y A D S V K G R F A V S R D Y A E N
 ACGGTGTATCTGCAAATGAACAGCCTGAAACCTGAGGACACGGCCGTTTATTACTGTAAC
 T V Y L Q M N S L K P E D T A V Y Y C N
 ACAAGGGCCTACTGGGGCCAGGGGACCCAGGTCACCGTCTCCTCAGCGCACCACAGCGAA
 T R A Y W G Q G T Q V T V S S A H H S E
 GACCCAGCTCCGCGGCCGCCCATCACCATCACCATCACGGGGCCGCAGAACAAAACTC
 D P S S A A A H H H H H H G A A E Q K L
 ATCTCAGAAGAGGATCTGAATTCTGAGAAAGATGAGCTATGACCAATTG
 I S E E D L N S E K D E L *MunI*

Fig.16.




Fig.17A.




Fig.17B.




Fig.18.

```

      NcoI          PstI
      -----
1  ccattggaggt gcagctgcag gaggtcagggg gaggattggt gcaggctggg
   >>.....HCV33.....>
      m e v q l q e s g g g l v q a g

51  ggctctctga gactctcctg tgcagcctcg ggaacgcgcca ccagtgggtca
   >.....HCV33.....>
      g s l r l s c a a s g r a t s g

101  tggtcactat ggtatgggct ggttcgcgcca ggttcaggg aaggagcgtg
   >.....HCV33.....>
      h g h y g m g w f r q v p g k e r

151  agtttgtcgc agctattagg tggagtggta aagagacatg gtataaagac
   >.....HCV33.....>
      e f v a a i r w s g k e t w y k d

201  tccgtgaagg gccgattcac catctccaga gataacgcca agactacggt
   >.....HCV33.....>
      s v k g r f t i s r d n a k t t

251  ttatctgcaa atgaacagcc tgaaacctga agatacggcc gtttattatt
   >.....HCV33.....>
      v y l q m n s l k p e d t a v y y

301  gtgcgcctcg accgggtccgc gtggatgata ttccctgcc ggttgggttt
   >.....HCV33.....>
      c a a r p v r v d d i s l p v g f
                        BstEII
                        -----
351  gactactggg gccaggggac ccaggtcacc gtctcctcag aaccaagac
   >.....HCV33.....>>>>..Hinge..>
      d y w g q g t q v t v s s e p k

401  accaaaacca caaccacaac cacaaccaca accacaaccc aatcctacaa
   >.....Hinge.....>
      t p k p q p q p q p q p q p n p t

451  cagaatccaa gtgtcccaaa tgtccagccc ctgagctcct gggagggccc
   >.....Hinge.....>>>>.....CH2.....>
      t e s k c p k c p a p e l l g g p

501  tcagtcttca tcttcccccc gaaacccaag gacgtctctt ccatttctgg
   >.....CH2.....>
      s v f i f p p k p k d v l s i s

551  gaggcccgag gtcacgtgcg ttgtggtaga cgtgggccag gaagaccccg
   >.....CH2.....>
      g r p e v t c v v v d v g q e d p

601  aggtcagttt caactgggtac attgatggcg cagaggtgcg aacggccaac
   >.....CH2.....>
      e v s f n w y i d g a e v r t a n

651  acgaggccaa aagaggaaca gttcaacagc acgtaccgcg tggtcagcgt
   >.....CH2.....>

```

Fig.18 (Cont).

```

      t r p k e e q f n s t y r v v s
701 cctgcccatac cagcaccagg actggctgac ggggaaagag ttcaaagtca
    >.....CH2.....>
    v l p i q h q d w l t g k e f k c
      HincII
      -----
751 aggtcaacaa caaagctctc ccggccccc tcgagaagac catctccaag
    >.....CH2.....>
    k v n n k a l p a p i e k t i s k

801 gccaaagggc agaccggga gccgcaggtg tacgccttg cccacacgg
    >...>>>.....CH3.....>
    a k g q t r e p q v y a l a p h

851 ggaagagctg gccaaaggaca ccgtgagcgt aacctgcctg gtcaaaggct
    >.....CH3.....>
    r e e l a k d t v s v t c l v k g

901 tctacccacc tgatatcaac gttgagtggc agaggaacgg tcagccggag
    >.....CH3.....>
    f y p p d i n v e w q r n g q p e

951 tcagagggca cctacgccac cagccaccc cagctggaca acgacgggac
    >.....CH3.....>

      s e g t y a t t p p q l d n d g

1001 ctacttcctc tacagcaagc tctcggtggg aaagaacacg tggcagcggg
    >.....CH3.....>
    t y f l y s k l s v g k n t w q r

1051 gagaaacctt cacctgtgtg gtgatgcaag aggccttgca caaccactac
    >.....CH3.....>
    g e t f t c v v m h e a l h n h y
                                EcoRI
                                -----
1101 acccagaaat ccatacccca gtcttcgggt aaataataag aattcgagct
    >.....CH3.....>>
    t q k s i t q s s g k

1151 cgaa

```

Fig.19.

```

      NcoI          PstI
      -----
1  ccatggagggt gcagctgcag gagtcagggg gaggattggt gcaggctggg
   >>.....HCV33.....>
      m e v q l q e s g g g l v q a g

51  ggctctctga gactctcctg tgcagcctcg ggaagcgcca ccagtgggtca
   >.....HCV33.....>
      g s l r l s c a a s g r a t s g

101  tggtcactat ggtatgggct ggttccgcca ggttccaggg aaggagcgtg
   >.....HCV33.....>
      h g h y g m g w f r q v p g k e r

151  agtttctctg agctattagg tggagtggta aagagacatg gtataaagac
   >.....HCV33.....>
      e f v a a i r w s g k e t w y k d

201  tccgtgaagg gccgattcac catctccaga gataacgcca agactacggt
   >.....HCV33.....>
      s v k g r f t i s r d n a k t t

251  ttatctgcaa atgaacagcc tgaaacctga agatacggcc gtttattatt
   >.....HCV33.....>
      v y l q m n s l k p e d t a v y y

301  gtgccgctcg accggctcgc gtggatgata ttccctgccc ggttggggtt
   >.....HCV33.....>
      c a a r p v r v d d i s l p v g f
                        BstEII
                        -----
351  gactactggg gccaggggac ccagggtcac gtctcctcag aacccaagac
   >.....HCV33.....>>>>..Hinge..>
      d y w g q g t q v t v s s e p k

401  accaaaacca caaccacaac cacaaccaca accacaaccc aatcctacaa
   >.....Hinge.....>
      t p k p q p q p q p q p q p n p t

451  cagaatccaa gtgtcccaaa tgtccagccc ctgagctcct gggagggccc
   >.....Hinge.....>>>>.....CH2.....>
      t e s k c p k c p a p e l l g g p

501  tcagtcttca tcttcccccc gaaacccaag gacgtcctct ccatttctgg
   >.....CH2.....>
      s v f i f p p k p k d v l s i s

551  gaggcccgag gtcacgtgcg ttgtggtaga cgtgggcccag gaagaccccg
   >.....CH2.....>
      g r p e v t c v v v d v g q e d p

601  aggtcagttt caactgggtac attgatggcg cagaggtgcg aacggccaac
   >.....CH2.....>
      e v s f n w y i d g a e v r t a n

651  acgaggccaa aagaggaaca gttcaacagc acgtaccgcg tggtcagcgt
   >.....CH2.....>

```

Fig.19 (Cont).

```

      t r p k e e q f n s t y r v v s
701 cctgcccatc cagcaccagg actggctgac ggggaaagag ttcaaatgca
    >.....CH2.....>
    v l p i q h q d w l t g k e f k c
      HincII
      -----
751 aggtcaacaa caaagctctc ccggccccc tcgagaagac catctccaag
    >.....CH2.....>
    k v n n k a l p a p i e k t i s k
801 gccaaagggc agaccggga gccgcaggtg tacgccttg cccacacgg
    >...>>>.....CH3.....>
    a k g q t r e p q v y a l a p h
851 ggaagagctg gccaaaggaca ccgtgagcgt aacctgcctg gtcaaaggct
    >.....CH3.....>
    r e e l a k d t v s v t c l v k g
901 tctaccacc tgatatcaac gttgagtggc agaggaacgg tcagccggag
    >.....CH3.....>
    f y p p d i n v e w q r n g q p e
951 tcagagggca cctacgccac caccgccccc cagctggaca acgacgggac
    >.....CH3.....>
      s e g t y a t t p p q l d n d g
1001 ctacttcttc tacagcaagc tctcgggtgg aaagaacacg tggcagcggg
    >.....CH3.....>
    t y f l y s k l s v g k n t w q r
1051 gagaaacctt cacctgtgtg gtgatgcacg aggccttgca caaccactac
    >.....CH3.....>
    g e t f t c v v m h e a l h n h y
1101 acccagaaat ccatcaccca gtcttcgggt aaatctgaga aagatgagct
    >.....CH3.....>>>.....SEKDEL.....>
    t q k s i t q s s g k s e k d e
      EcoRI
      -----
1151 ataataagaa ttcgagctcg aa
    >
    l

```

Fig.20.




Fig.21.




Fig.22.




Fig.23.




Fig.24.




Fig.25.




Fig.26.





Fig.27.



80

Fig.29.

